

Fortegnelse, kontroldokument og brugervejledning kategori A – MFAG A1

Inventory, Control Document and User Instructions Category A – MFAG A1



Fortegnelse, kontroldokument og brugervejledning over lægemiddelstoffer og medicinsk udstyr kategori A - MFAG A1

Jf. Meddelelser fra Søfartsstyrelsen A, Kapitel IX B

Inventory, Control Document, User Instructions on Active Pharmaceutical Ingredients and Medical Equipment Category A – MFAG A1

Cf. Danish Maritime Authority Notice A, chapter IX B

July 2025

ISBN: 978-87-7454-211-7

Identifikation af skib / Identification of ship

Skibsnavn / <i>Ship Name:</i>	
Kaldesignal / <i>Call Signal:</i>	
IMO nummer / <i>IMO number:</i>	
Hjemmehavn / <i>Port of Reg.:</i>	

Medicin certifikat / *Medicine Certificate*

Dette dokument anses for at udgøre skibets medicin certifikat.

Det attesteres herved, at medicin og medicinsk udstyr er efterset og fundet i overensstemmelse med
Den Danske Søfartsstyrelses krav jf. Meddelelser A Kapitel IX B.

This document is considered to represent the ships certificate of medicine.

*It certifies that medicine and medical equipment has been inspected and found in compliance with
the Danish Maritime Authority's requirement cf. Notes A chapter IX B.*

Skibsnavn / *Ship Name*:

Kaldesignal / *Call Signal*:

IMO nummer / *IMO number*:

Hjemmehavn / *Port of Reg.*:

Dato / <i>Date</i>	Rank / <i>Rank</i>	Navne / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / Master		
Næste eftersyn / <i>Next Inspection</i>			
Dato / <i>Date</i>	Rank / <i>Rank</i>	Navne / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / Master		

Print denne side / *Print this page*

Certifikat for genoplivningsudstyr (A.1)

Certificate resuscitation equipment (A.1)

Det attesteres herved at det **mobile genoplivningsudstyr (A.1)** er efterset og fundet i overensstemmelse med Den Danske Søfartsstyrelses krav jf. Meddelelser A Kapitel IX B.

It certifies that the **mobile resuscitation equipment (A.1)** has been inspected and found in compliance with the Danish Maritime Authority's requirement cf. Notes A chapter IX B.

Skibsnavn / *Ship Name*:

Kaldesignal / *Call Signal*:

IMO nummer / *IMO Number*:

Hjemmehavn / *Port of Reg.*:

Dato / <i>Date</i>	Rank / <i>Rank</i>	Navn / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		
Næste eftersyn / <i>Next Inspection</i>			
Dato / <i>Date</i>	Rank / <i>Rank</i>	Navne / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		

Print denne side / *Print this page*

Indhold / Contents

Brugervejledning:	5
<i>User Instruction:</i>	5
Anvendte forkortelser	9
<i>Used abbreviations</i>	9
Brugervejledning Lægemiddelstoffer	10
<i>User instruction Activ Pharmaceutical</i>	10
Fortegnelse og kontroldokument	64
<i>Inventory and Control Document</i>	64
Besætningens medbragte medicin / Crew member's private medicine	71
Fortegnelse og Kontrolldokument	73
<i>Inventory and Control document</i>	73
Labels	82
INDEX	87
Rekvistition lægemiddelstoffer	92
<i>Requisition Active Phamaceutecal</i>	92
Mobilt genoplivnings udstyr / Akut taske	104
<i>Mobile resuscitation kit / Emergency bag</i>	104
Ændringer / Changes	108

De Danske Medicinkister

Medicinkisten angiver medicin/medicinsk udstyr, som Radio Medical Danmark og Søfartsstyrelsen har foretrukket indenfor de rammer som EU-direktiv 92/29/EØF og den Danske lægemiddelstyrelse har givet.

Guide til:

Brugervejledning:

Første kolonne beskriver de indikationer, lægemiddelstoffet er beregnet til. De samme indikationer kan slås op i indekset bagerst i hæftet. Første kolonne beskriver også om et lægemiddelstof skal anvendes sammen med andre lægemiddelstoffer. I sådanne tilfælde er der lavet henvisninger til de relevante lægemiddelstoffer.

I samme kolonne gøres opmærksom på, om lægemiddelstoffet udgør en del af MFAG (Medical First Aid Guide), der refereres i så fald til den relevante MFAG tabel.

MFAG lægemiddelstofferne er defineret i IMDG (International Maritime Dangerous Goods) koden, supplementdelen, seneste version og indarbejdet i denne version af den danske "Fortegnelse, kontrol dokument og brugervejledning". kontrol dokument og brugervejledning".

Bemærkning: Sejlads med farligt gods jævnfør IBC koden, INF koden og IGS koden er ikke indarbejdet i denne publikation. Der henvises til disse publikationer.

Anden kolonne oplyser, Lægemiddelstoffets hoved- og undergruppenummer.

Tredje og fjerde kolonne indeholder rækker med oplysninger om:

- **Lægemiddelstoffet:** Beskriver det aktive stof i lægemidlet. Som et eksempel kan nævnes 3.1 Paracetamol. Det forhandles over hele jorden under mange forskellige handelsnavne. Derfor kan man uanset forskellige handelsnavne, altid finde lægemidlet, hvis man kender lægemiddelstoffet.
- **Form:** Beskriver om lægemiddelstoffet skal findes som kapsel, tablet, suppositorier, injektionsvæske, infusionsvæske, creme, salve, afdelte pulver, mund spray, brusetablet, klyksma, oral opløsning eller inhalation. Form beskriver også styrken af det enkelte lægemiddelstof pr ml eller per tablet.
- **Virkning** beskriver den forventende virkning af det enkelte lægemiddel.
- **Dosering** beskriver den anbefalede dosis ved de tilstande, der er beskrevet i kolonnen indikation. Her beskrives ligeledes lægemidlets form og administrationsmåde.
Bemærkning: Radio Medical kan ordinere andre doser og administrations måder.
- **Bivirkninger** nævner de mest almindelige bivirkninger beskrevet i Dansk Lægemiddelinformation, **www.medicin.dk**
- **Holdbarhed** beskriver holdbarhedstiden fra produktions datoen. Holdbarhedstiden beskrevet i denne fortegnelse er fremkommet ud fra det danske handelsnavn med længst holdbarhed.
Bemærkning: Der tages forbehold for at holdbarhedstiden kan ændres.
- **Opbevaring** beskriver om der er særlige krav til opbevaring. Lægemidler skal generelt opbevares tørt og beskyttet mod lys. De fleste lægemidler tåler "almindelig temperatur" defineret som normal stuetemperatur op til 22-24° C.

The Danish Medicine Chest's

The medicine chest indicates the Active Pharmaceutical Ingredients / medical equipment that Radio Medical Denmark and Danish Maritime Authority have preferred within the framework of the EU Directive 92/29 / EEC and the Danish Medicines Agency.

Guide to:

User Instruction:

First column describes the indication the Active Pharmaceutical Ingredients is intended for. The same indications can be looked up in the index at the back of the booklet. First column also describes whether an Active Pharmaceutical Ingredient should be used together with other Active Pharmaceutical Ingredients. In such cases, a reference is made to the relevant Active Pharmaceutical Ingredient. In the same column it is noted if the Active Pharmaceutical Ingredients is part of MFAG (Medical First Aid Guide), and reference to the relevant MFAG table is made. MFAG Active Pharmaceutical Ingredients s are defined in the IMDG (International Maritime Dangerous Goods) code, supplement component, latest edition, and incorporated in this version of the Danish "Inventory, control document and user manual".

Note: Sailing with dangerous goods as per the IBC code, INF code and the IGC code are not included in this publication. Please refer to these publications.

Second column describes the Active Pharmaceutical Ingredient main and subgroup numbers.

Third and fourth column contains a number of information about:

- **The Active Pharmaceutical Ingredients:** describe the Active Pharmaceutical Ingredients in the medicine. An example is 3.1 Paracetamol, which is sold all over the world under many different trade names. Regardless of different trade names it is always possible to locate the medicine, if you know the Active Pharmaceutical Ingredients.
- **Form:** describes if the Active Pharmaceutical Ingredients is available as capsules, tablets, suppositories, injection liquid, infusion liquid, lotion, ointment, divisional powder, mouth spray, effervescent tablets, enema, oral solution or inhalation. Form also describes the strength of each active pharmaceutical ingredient per ml or per tablet.
- **Effect** describes the expected effect of each Active Pharmaceutical Ingredients.
- **Dosage** describes the recommended dose for the condition described in the column indication. The form of the Active Pharmaceutical Ingredients as well as the usual administration is mentioned.
Note: Radio Medical may prescribe different doses and ways of administration.
- **Side effects** describes the most common side effects mentioned in the Danish Drug Information, **www.medicin.dk**. There is currently no English version.
- **Durability** describes the time of durability from date of production. The durability mentioned in this record is derived from the Danish trade name with the longest durability.
Note: Durability may be subject to change.
- **Storage** describes if special storage precautions are needed. The Active Pharmaceutical Ingredients should generally be kept dry and protected from light. Most medicaments tolerate "normal temperature" defined as normal room temperature up to 22-24° C.

- **Anmærkninger** beskriver, om der bør søges samråd med Radio Medical Danmark inden anvendelse af lægemiddelstoffet. Desuden beskrives, om der er specielle forholdsregler til det specifikke lægemiddelstof.

Advarsel: Lægemiddelstoffer, der har været opbevaret forkert, eller er blevet for gamle, kan være sundhedsskadelige.

Guide til:

Fortegnelse og kontroldokument over lægemiddelstoffer

Publikationen indeholder en fortegnelse over de lægemiddelstoffer, der skal være ombord.

Rækkene

Lægemiddelstofferne er organiseret i hovedgrupper efter deres hovedanvendelsesområde. Eksempelvis er alle smertestillende lægemiddelstoffer organiseret i gruppe 3.

Kolonnerne

Første kolonne beskriver hvert enkelt lægemiddelstoffs nummer. Eksempelvis har morfin nummeret 3.4, hvor tallet før punktum referer til hovedgruppen, og tallet efter refererer til det specifikke lægemiddelstof. Lægemiddelstofferne i skibets hospital skal så vidt muligt organiseres efter dette nummersystem

Anden kolonne beskriver ATC koden, et klassifikations system for medicin. Se mere på www.whocc.no

Tredje kolonne beskriver lægemiddelstoffet.

Bemærkning: Efter lægemiddelstoffet kan være anført et "Δ", dersom produktet er sløvende, og et "K", dersom produktet skal opbevares i køleskab.

Fjerde kolonne beskriver under "Adm.form", om lægemiddelstoffet skal findes som kapsel, tablet, suppositorier, injektionsvæske, infusionsvæske, creme, salve, afdelte pulver, mund spray, brusetablet, klysma, oral opløsning eller inhalation.

Femte kolonne beskriver styrken pr tablet eller pr. ml.

Sjette kolonne beskriver mindste mængde, skibet skal medbringe. Skibet må gerne medbringe større mængder end det beskrevne.

Syvende kolonne MFAG A1 beskriver det supplement, skibet skal medbringe, dersom det sejler med A-fortegnelse OG har en evakueringstid på 24 timer eller mere.

Bemærkning: Kolonnerne mærket MFAG beskriver de minimumsmængder af supplerende lægemidler som skibet skal medbringe, dersom skibet sejler med farligt gods.

- **Remarks** describe whether it is recommended to seek advice from Radio Medical Denmark before using the Active Pharmaceutical Ingredients. In addition, precautions needed for the specific Active Pharmaceutical Ingredients are described.

Warning: Active Pharmaceutical Ingredients which have been stored improperly or are too old can be harmful to health.

Guide to:

Inventory and Control Document of Active Pharmaceutical Ingredients.

The publication contains an inventory of mandatory Active Pharmaceutical Ingredients on board.

The rows

The Active Pharmaceutical Ingredients are organized into main groups according to field of application. As an example all analgesics are organized in group 3.

The columns

First column describes each individual Active Pharmaceutical Ingredient by number. For example, morphine number 3.4, where the number before the dot refers to the main group and the number after the dot refers to the specific Active Pharmaceutical Ingredients. The Active Pharmaceutical Ingredients in the ship's hospital must as far as possible be organized after this numbering system.

Second column describe the ATC code, a classification system on medicine. See more on www.whocc.no.

Third column describes the Active Pharmaceutical Ingredients

Note: The Active Pharmaceutical Ingredients can be marked with a "Δ" if the product is sedative and a "R" if the product should be stored in a refrigerator

Fourth column describes under "Adm.form" whether the Active Pharmaceutical Ingredients should be available as capsules, tablets, suppositories injection liquid, infusion liquid, lotion, ointment, divisional powder, mouth spray, effervescent tablets, enema, oral solution or inhalation.

Fifth column describes the strength per tablet or per ml.

Sixth column describes the minimum amount the vessel must carry. The vessel may carry greater amounts than described.

Seventh column MFAG A1 describes the supplement the ship must bring if it sails with A-record AND has an evacuation time on 24 hours or more.

Note: The columns marked MFAG describes the minimum quantities of additional (supplement) Active Pharmaceutical Ingredients the ship must bring, if the ship is carrying dangerous goods.

Besætningens medbragte medicin

Besætningen har ret til at medbringe egen medicin under den forudsætning, at de til stadighed har en gyldig helbredsundersøgelse for søfarende og fiskere. Medbragte lægemidler skal oplyses til skibsføreren eller sygdomsbehandleren.

Lægemidlet skal indskrives med følgende oplysninger:

- Lægemidlets navn
- Navn på besætningsmedlem
- Mængde og antal
- Daglig dosis
- Dato ind
- Eventuel kvittering for modtagelsen. (Kun hvis lægemidlet opbevares på sygelukafet)
- Dato ud
- Kvittering for modtagelsen (Kun hvis lægemidlet har været opbevaret på sygelukafet)

Guide til:

Fortegnelse og kontroldokument over medicinsk udstyr

Publikationen indeholder en fortegnelse over det medicinske udstyr, der skal være om bord.

Rækkerne

Det medicinske udstyr er organiseret i hovedgrupper efter deres hovedanvendelsesområde. Eksempelvis er alle forbindinger og plastre organiseret i gruppe B.

Første kolonne beskriver det enkelte medicinske udstyrs bogstav og nummer. Eksempelvis har "Elastisk gazebind, ca. 4 cm x 4 m", kombinationen nummeret B.9, hvor bogstavet før punktum referer til hovedgruppen, og tallet efter referer til det specifikke medicinske udstyr.

Anden kolonne beskriver det specifikke medicinske udstyr.

Tredje kolonne beskriver den mindste mængde, skibet skal medbringe. Skibet må gerne medbringe større mængder end det beskrevne.

Fjerde kolonne MFAG A1 beskriver det supplement, skibet skal medbringe, dersom det sejler med A-fortegnelse OG har en evakuerings tid på 24 timer eller mere.

Bemærkning: Kolonnerne mærket MFAG beskriver de minimumsmængder af supplerende medicinsk udstyr som skibet skal medbringe, dersom skibet sejler med farligt gods

Det mobile genoplivningsudstyr A.1

Søfartsstyrelsen har fastsat regler om ilt og det mobile genoplivningsudstyr i gældende Meddelelser A kapitel IX B. Man kan derudover pakke førstehjælpsudstyr og lægemidler til førstehjælp sammen med det mobile genoplivningsudstyr. Der er udarbejdet forslag til pakning af et mobilt genoplivningsudstyr på www.soefartsstyrelsen.dk.

Etiketter

Publikationen indeholder en kopi af de etiketter, der skal være på lægemiddelstofferne ved leveringen. Kopien kan anvendes som kontrol af, at lægemidlet er korrekt og påført den korrekte etiket. Er det ikke tilfældet, bør skibsføreren rettlede sin leverandør.

Ved køb af lægemiddelstoffer, kan en kopi af etiketterne afleveres ved leverandøren. Derefter skal leverandøren, før udlevering, påføre de korrekte etiketter,

Indeks

Indekset henviser til lægemiddelstoffernes brugervejledning. I indekset kan du søge på visse sygdomme, symptomer og alle lægemiddelstoffer.

Private medicine of crewmembers

The crew has a right to bring their own medicine, provided they possess a valid health certificate for seafarers and fishermen at all time. The master or the person in charge of medical care has to be informed about medicine brought along.

The medicine must be registered with the following information:

- Name of the medicine
- Name of crewmember
- Amount
- Daily dosage
- Date in
- Possible receipt for received medicine.
(Only if the medicine is stored at the sickbay)
- Date out
- Sign when medicine is returned
(Only if the medicine has been stored in the sickbay)

Guide to:

Inventory and Control Document medical equipment

The publication contains an inventory of the mandatory medical equipment onboard

The rows

The medical equipment is organized into main groups according to their main field of application. For example all dressings and plasters are organized in group B.

First column describes each individual medical equipment by number. The "elastic gauze", approx. 4 cm x 4 m, carries the combination B.9. The letters before the dot refers to the main group, and the number after the dot refers to the specific medical equipment.

Second column describes the specific medical equipment.

Third column describes the minimum amount the vessel must carry. The vessel may carry larger quantities than described.

Fourth column MFAG A1 describes the supplement the ship must bring if it sails with A-record AND has an evacuation time on 24 hours or more.

Note: The columns marked MFAG describes the minimum quantities of additional (supplement) medical equipment the ship must bring, if the ship is carrying dangerous goods.

The Mobile Resuscitation Equipment A.1

The Danish Maritime Authority has established rules for oxygen and the mobile resuscitation equipment in the current Notice A chapter IX B. In addition you may pack equipment and medicine suited for first aid with the resuscitation equipment. Suggestions for the list may be found on www.dma.dk.

Labels

The publication contains copies of the labels that must be on each active pharmaceutical ingredient, at the time of delivery. The copy may be used as verification of proper product and label. If labelling is incorrect, the master should guide his supplier.

When purchasing Active Pharmaceutical Ingredients a copy of the label may be delivered to the supplier. Then the supplier applies the correct labels before delivery.

Index

The index refers to the user instruction of active pharmaceutical ingredients. You can search for diseases, symptoms and Active Pharmaceutical Ingredients in the index.

Rekvisition

I publikationen findes en bestillingsliste, hvor du blot udfylder de ønskede mængder af lægemiddelstof og medicinsk udstyr.

Medicinkiste- / Mobilt genoplivningsudstyr certifikat.

Mindst en gang årligt skal skibets lægemiddelstoffer og medicinske udstyr gennemgås.

Det mobile genoplivnings udstyr skal kontrolleres hver 3 måned. Ved gennemgangen kontrolleres mængden, opbevaringsforhold, udløbsdato og etiketter. Nåle, kanyler og metalinstrumenter kontrolleres for rust. Kontroller, at emballagen er intakt. Vær opmærksom på, at lægemiddelstof og medicinsk udstyr skal opbevares tørt og mørkt, og ikke må opbevares i nærheden af klor.

Dokumentation af 3 måneders og årlig inspektion.

På side 2 og 3 er Medicinkiste – og mobilt genoplivningsudstyrs certifikat, som også kan anvendes til dokumentation af inspektionerne. Siderne udskrives og underskrives af skibsføreren.

Dispensation for andet lægemiddelstof

Det er ikke til alle tider, og i alle lande muligt at indkøbe det foretrukne lægemiddelstof. Den danske Søfartsstyrelse og Radio Medical Danmark har derfor udarbejdet en liste med godkendte alternativer. Hvis det er muligt at indkøbe lægemiddelstoffer fra listen med godkendte alternativer med den korrekte koncentration, og indgiftsmåde, er det ikke nødvendig at ansøge om dispensation.

Hvis apotek/forhandler tilbyder lægemiddelstof, som ikke findes på nogen liste, skal der søges om dispensation for anden medicin. Dette gøres på Søfartsstyrelsen hjemmeside, hvor der er en procedure for "Dispensation for anden medicin".

Listen over godkendte alternative, og dertil hørende labels og brugervejledninger kan downloades fra www.sofartsstyrelsen.dk.

Listen med godkendte alternativer findes i Excel format.

Hjælp

Er der noget du er i tvivl om, står Søfartsstyrelsens "Center for det Maritime Sundhedsvæsen" til din rådighed. Adressen er:

CMS, Vestervejen 1, 6720 Fanø
Telefon: +45 7219 6004
Email: cms@dma.dk

Requisition

In the publication is a requisition form available. You simply fill in the required quantities of the active pharmaceutical ingredients and medical equipment.

Medicine chest- / Mobile resuscitation equipment certificate.

At least once a year the ship's Active Pharmaceutical Ingredients and medical equipment must be reviewed. The mobile resuscitation equipment every 3 month.

When reviewing the quantities, storage conditions, expiry date and labels are controlled. Needles, cannulas and metal instruments are checked for rust. Check if the packaging is intact. Be aware that the Active Pharmaceutical Ingredients and equipment must be stored dry and dark, and not near chlorine.

Documentation of 3 months and annual inspection.

On page 2 and 3 is Medicine chest- and mobile resuscitation equipment's certificate's, it can be used to document the inspections. The page's has to be printed and the Master has to sign them.

Exemption for other active pharmaceutical ingredients

It is not at all times and in all countries possible to purchase the preferred Active Pharmaceutical Ingredients. The Danish Maritime Authority and Radio Medical Denmark have therefore drawn up a list of approved alternatives. If it is possible to purchase Active Pharmaceutical Ingredients from the list of approved alternatives with the correct concentration and method of administration, it is not necessary to apply for exemption.

If the pharmacy / retailer offer Active Pharmaceutical Ingredients that is not on the inventory- or alternative list, you must apply for an exemption. This is done on the DMA website, where there is a procedure for "exemption of medicine".

The list of approved alternative, and associated labels and user instructions can be downloaded from www.dma.dk. The list of approved alternatives is available in Spreadsheet format.

Help

If you have any questions, the Danish Maritime Authority's "Centre of Maritime Health Service" is at your disposal. The address is:

CMS, Vestervejen 1, 6720 Fanø
Telephone: +45 7219 6004
E-mail: cms@dma.dk

Anvendte forkortelser m.v. i Fortegnelse og Brugervejledning

afd. Pul.	=	afdelte pulvere
gel	=	gel til påsmøring på hud
htgl.	=	hætteglas
i. m.	=	intramuskulært = i musklen
inf. Væs.	=	infusionsvæske (til intravenøs brug)
inh. Pul.	=	inhalationspulver
inj. væske	=	injektionsvæske (til i.m. eller s.c.injektion)
i.v.	=	intravenøst = i en vene
"K"	=	opbevares i køleskab
klysma	=	klysma til anvendelse i endetarmen
linim.	=	liniment
gas	=	luftformig gasart til medicinsk anvendelse
mixtur	=	væske som drikkes
næsedr.	=	næsedråber
opløsn.	=	opløsning
s.c.	=	subcutant = under huden
supp.	=	suppositorier = stikpille til brug i endetarmen
Δ	=	sløvende medicin, kan påvirke evnen til maskinbetjening og vagttjeneste
(π)	=	Receptpligtig medicin

Used abbreviations etc. In the Inventory and User instruction

d.p.	=	divisional powder
gel	=	gel for application on the skin
c.v	=	cappedvials
i.m.	=	intramuscular = in the muscle
inf. fl.	=	infusion fluid (for intravenous use)
inh. p.	=	inhalation powder
inj. subst.	=	injection substance (for reconstitution before use)
inj. liquid	=	injection fluid (for i.m. or s.c. injection)
i.v.	=	intravenous = in a vein
"R"	=	stored in refrigerator
enema	=	enema for use in the rectum
linim.	=	liniment
gas.	=	gas for medical use
mixture	=	drinkable liquid
nose d.	=	nose drops
diss.	=	dissolution
s.c.	=	subcutaneous = under the skin
supp.	=	suppositories = suppository for rectal use
Δ	=	sedative medicine, may effect the ability to operate machinery and bridge duty
(π)	=	Prescription drugs

Brugervejledning Lægemiddelstoffer

**User instruction Active Pharmaceutical
Ingredients**

Grp. 1 Ilt / Oxygen

Bevidstløs uden vejrtrækning Bevidstløs med vejrtrækning Medtaget almen tilstand Slag mod kraniet Større kvæstelser Vejrtrækningsbesvær MFAG tabel 2, 3, 5, 9, 11	1.1+1.3	Lægemiddelstof:	Medicinsk ilt
		Form:	Luft på trykflasker.
		Virkning:	Tilfører ilt til cellerne via de røde blodlegemer.
		Dosering:	Bevidstløse uden vejrtrækning: Manuel ventilation med 15 liter per minut. Bevidstløse med vejrtrækning og øvrige tilstande: Maske med 9 liter/min.
		Bivirkninger:	Ilt er ufarligt til medicinsk brug.
		Holdbarhed:	Flaske: 3 år fra produktionsdato (Conoxia®, Linde Gas Therapeutics AGA).
		Opbevaring	Holdes væk fra brændbare gasser og stoffer. Flasker holdes under 50°C.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Unconscious without breathing Unconscious with breathing Weak general condition Blows to the skull Severe injuries Breathing difficulties MFAG table 2, 3, 5, 9, 11	1.1+1.3	Active Pharmaceutical Ingredients:	Medical oxygen
		Form:	Gas in pressurized bottles.
		Effect:	Induces oxygen to the cells via the red blood cells.
		Dosage:	Unconscious without breathing: Manual ventilation, 15 litres per minute. (A.1). Unconscious with breathing and other states: Oxygen mask, 9 litres per minute.
		Side-effects:	Oxygen is not dangerous when used for medicinal purposes.
		Validity:	Bottles 3 years from productions date (Conoxia®, Linde Gas Therapeutics AGA).
		Storage:	Keep away from flammable gases and substances. Keep bottle under 50°C.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 2 Lokalbedøvelse / Local Anaesthetics

Lokalbedøvelse Syning af sår	2.1	Lægemiddelstof:	Lidocain
		Form:	Injektionsvæske 10 mg/ml til subcutan injektion.
		Virkning:	Lokalbedøver på injektionsstedet. Virker efter 1 - 2 minutter. Virker i 2 - 3 timer.
		Dosering:	Mængden afhænger af sårets størrelse. Normalt anvendes 20 - 50 mg. Maksimaldosis er 20 ml svarende til 200 mg. Se Søfartsstyrelsens Lægebog.
		Bivirkninger:	Allergi er sjælden. Ved utilsigtet injektion i blodbanen kan der opstå: 1. føleforstyrrelser, svimmelhed, uklarhed og øresusen. 2. synsforstyrrelser, muskelrysten og kramper. 3. blodtryksfald og langsom puls.
		Holdbarhed:	3 år fra produktionsdato (Xylocain®, AstraZeneca).
		Opbevaring:	Almindelig temperatur. Må ikke fryses.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Lokalbedøvelse skal ske i et rum med adgang til adrenalin (4.1) og genoplivningsudstyr (A.1). Risiko for alvorlige bivirkninger, hvis lidocain injiceres i blodbanen.

Local anaesthetics Suture of wounds	2.1	Active Pharmaceutical Ingredients:	Lidocaine
		Form:	Injection liquid, 10 mg/ml for subcutaneous injection.
		Effect:	Anaesthetises the place of injection. Takes effect after 1-2 minutes. Effective for 2-3 hours.
		Dosage:	The amount depends on the size of the wound. Normally, 20-50 mg is used. The maximum dose is 20 ml, equivalent to 200 mg. See the Danish Maritime Authority's Medical Manual.
		Side-effects:	Allergy seldom occurs. In case of an unintended injection into the bloodstream, the following side-effects may occur: 1. affective disturbances, dizziness, confusion, and buzzing in the ears. 2. visual disturbances, muscle contractions, and convulsions. 3. falling blood pressure and slow pulse.
		Validity:	3 years from the date of production (Xylocain®, AstraZeneca).
		Storage:	Normal temperature. Not to be frozen.
Lokalbedøvelse Kløestillende	2.2	Remarks:	Consult <i>Radio Medical</i> . Local anaesthetics must be administered in a room with access to adrenaline (4.1) and resuscitation equipment (A.1). Risk of serious side-effects if lidocaine is injected directly into the bloodstream.
		Lægemiddelstof:	Lidocain
		Form:	Gel, 20 mg/g (2%).
		Virkning:	Lokalbedøver hud og sårrende. Kløestillende.
		Dosering:	Anvendes som lokalbedøvelse ved smertende sår eller smertefulde procedurer, fx ved rengøring af brandsår eller større og forurenede sår. Et tykt lag gel påsmøres huden, der herefter afdækkes lufttæt i ca. 15 minutter. Virker efter ca. 15 minutter. Virker i ca. 30 minutter.
		Bivirkninger:	Allergi er sjælden.
		Holdbarhed:	2 år fra produktionsdato (Xylocain®, AstraZeneca).
		Opbevaring:	Almindelig temperatur.
Local anaesthetics Relieves itching	2.2	Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
		Active Pharmaceutical Ingredients:	Lidocaine
		Form:	Gel, 20 mg/g (2%).
		Effect:	Anaesthetises the skin and the edges of wounds. Relieves itching.
		Dosage:	To be used as local anaesthetics in case of painful wounds or procedures, e.g. cleansing of burns or large, dirty wounds. A thick layer is applied to the skin, which is subsequently covered in an air-proof condition for about 15 minutes. Takes effect after about 15 minutes. Effective for about 30 minutes.
		Side-effects:	Allergy seldom occurs.
		Validity:	2 years from the date of production (Xylocain®, AstraZeneca).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 3 Smertestillende midler / Analgetics

Feber Svage smerter: Hovedpine Muskelsmerter Ledsmerter Rygsmarter MFAG tabeller: 7, 8, 13	3.1	Lægemiddelstof:	Paracetamol
		Form:	Tabletter à 500 mg.
		Virkning:	Febernedsættende og smertestillende.
		Dosering:	Voksne: 1 g 3 - 4 gange dagligt. Højst 4 g per døgn.
		Bivirkninger:	Doser over 15 - 20 g indebærer risiko for leverskade, leversvigt og død.
		Holdbarhed:	5 år fra produktionsdato (Pamol Nycomed DK ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Ingen.
Fever Weak pains: Headache Muscle pains Pain in joints Back pains MFAG tables: 7, 8, 13	3.1	Active Pharmaceutical Ingredients:	Paracetamol
		Form:	Tablets of 500 mg.
		Effect:	Pain relieving and reduces fever.
		Dosage:	Adults: 1 g three-four times daily. Not more than 4 g daily.
		Side-effects:	Doses of more than 15-20 g involve liver damage, liver failure, and death.
		Validity:	5 years from the date of production (Pamol Nycomed DK ApS).
		Storage:	Normal temperature.
		Remarks:	None.
Moderate smerter: Menstruationssmerter Ledsmerter Knoglesmerter Knoglebrud Muskelsmerter	3.2	Lægemiddelstof:	Ibuprofen
		Form:	Tabletter à 400 mg.
		Virkning:	Smertestillende og febernedsættende.
		Dosering:	Voksne: 400 mg 3 - 4 gange dagligt. Højst 1800 mg per døgn. Undtagelsesvist op til 2400 mg per døgn i kortere perioder (4 - 6 uger).
		Bivirkninger:	Ondt i maven.
		Holdbarhed:	5 år fra produktionsdato (Ibumetin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes ved: 1. allergi for acetylsalicylsyre. 2. astmapatienter. 3. patienter i blodfortyndende behandling. 4. patienter med symptomer på mavesår.

Moderate pains: Period pains Pain in joints Bone-ache Fractures Muscle pains	3.2	Active Pharmaceutical Ingredients:	Ibuprofen
		Form:	Tablets of 400 mg.
		Effect:	Pain relieving and reduces fever.
		Dosage:	Adults: 400 mg three-four times daily. Not more than 1800 mg daily. Exceptionally, up to 2400 mg daily for shorter periods of time (4-6 weeks).
		Side-effects:	Stomach ache.
		Validity:	5 years from the date of production (Ibumetin®, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> Not to be used in case of: 1. allergy to acetylsalicylic acid. 2. asthma patients. 3. patients in blood diluting treatment. 4. patients with symptoms of ulcer.
Stærke smerter: Store sårskader Kvæstelser Forbrændinger Knoglebrud Nyresten Galdesten Blodprop i hjertet (se førstehjælp under 9.1 Nitroglycerin) MFAG tabel 7, 8,10,13	3.4	Lægemiddelstof:	Morfin Δ
		Form:	Injektionsvæske 10 mg/ml, ampuller à 1 ml.
		Virkning:	Smertestillende. Vækker velvære, virker afslappende. Virker efter 5 – 15 min ved intramuskulær (i.m.) eller subcutan (sc.) injektion. Virker omgående ved intravenøs (i.v.) injektion. Virkningsvarighed: 3 -4 timer.
		Dosering:	Subcutant og intramuskulært (voksne). 10 mg. Kan gentages, hvis tilstrækkelig effekt udebliver. Intravenøst (voksne): 5 mg (halv ampul) injiceres langsomt. Kan gentages, hvis tilstrækkelig effekt udebliver.
		Bivirkninger:	Svækker vejtrækningen. Kvalme og opkastning. Hæmmer optagelsen af medikamenter fra mavesækken. Kan give forstoppelse og hæmme blæretømningen.
		Holdbarhed:	2 år fra produktionsdato, (Morfin, Skanderborg apotek).
		Opbevaring:	Almindelig temperatur. Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved overdosering kan vejtrækningen ophøre. Modgift: Naloxone (5.1). Denne bivirkning og morfins virkning i øvrigt forstærkes af andre beroligende stoffer som alkohol, sovemidler og lignende. Overdosering er især en risiko ved hurtig injektion. Overdosering sker sjældent ved almindelig dosering. Observer vejtrækningen og ventilere om nødvendigt med ilt (1.1) på ventilationspose og maske (A.1).

Severe pain Major wounds Injuries Burns Fractures Kidney stone Gall stone Coronary thrombosis (see first aid under 9.1 Glyceryl trinitrate) MFAG Tabel 7,8,10,13	3.4	Active Pharmaceutical Ingredients:	Morphine Δ
		Form:	Injection liquid, 10 mg/ml, ampoules of 1 ml.
		Effect:	Pain relieving. Gives the patient a feeling of well-being, relaxant. Takes effect after about 5-15 minutes in case of intramuscular or subcutaneous injection. Takes effect immediately in case of intravenous injection Effective for 3-4 hours.
		Dosage:	Subcutaneously and intramuscularly (adults): 10 mg. May be repeated if the effect is not sufficient. Intravenously (adults): 5 mg (1/2 ampoule) to be injected slowly. May be repeated if the effect is not sufficient.
		Side-effects:	Restricts breathing. Nausea and vomiting. Restricts the absorption of medicaments from the stomach. May cause constipation and restrict bladder emptying.
		Validity:	2 years from the date of production (Morphine, Skanderborg Pharmacy).
		Storage:	Normal temperature. To be protected from light.
		Remarks:	Consult <i>Radio Medical</i> . In case of an overdose, the patient may stop breathing. Antidote: Naloxone (5.1) This side-effect and the effect of morphine as drowsiness may be increased by the use of other sedatives, such as alcohol, sleeping medicine, ect. Risk of overdose primarily if the injection is given too quickly. Overdoses are rare in case of normal dosage. Observe the breathing and if necessary, ventilate by means of oxygen (1.1) ventilations bag and mask (A.1).
Inflammation i huden Gigt Smerter	3.7	Lægemiddelstof:	Diclofenac
		Form:	Gel 11,6 mg/g.
		Virkning:	Smertestillende.
		Dosering:	Smøres på huden i et tyndt lag 3-4 gange i døgnet.
		Bivirkninger:	Rødme, hududslæt, hudkløe, eksem, hududslæt ved kontakt med midlet.
		Holdbarhed:	3 år, fra produktionsdato (Voltaren® gel, Novatis Healthcare).
		Opbevaring:	Almindelig temperatur.
Inflammation of the skin Arthritis Pain.	3.7	Active Pharmaceutical Ingredients:	Diclofenac
		Form:	Ointment with 11,6 mg/g.
		Effect:	Analgesic.
		Dosage:	Apply a thin layer to the skin 3-4 times daily.
		Side-effects:	Redness, rash, itching, eczema, skin rashes when in contact with the product.
		Validity:	3 years from the date of production (Voltaren®gel, Novatis Healthcare A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Only to be used on intact skin. May not be used in the face.

Nyresten Galdesten Ledsmarter	3.8	Lægemiddelstof:	Diclofenac
		Form:	Suppositorier 50 mg.
		Virkning:	Smertestillende.
		Dosering:	50 -100 mg mg x 1 dagligt.
		Bivirkninger:	Nedsat appetit, sure opstød og halsbrand, kvalme, opkastning, mavesmerter, diarré, luft i maven. Hovedpine, svimmelhed. Hududslæt og hudkløe.
		Holdbarhed:	3 år fra produktionsdato, (Voltaren, 2care4).
		Opbevaring:	Almindelig temperatur - ikke over 30°C.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes ved: 1. allergi for acetylsalicylsyre. 2. astmapatienter. 3. patienter i blodfortyndende behandling. 4. patienter med symptomer på mavesår.
Kidney stone Gall stone Pain in joint	3.8	Active pharmaceutical ingredients:	Diclofenac
		Form:	Suppositorie 50 mg.
		Effect:	Analgesic.
		Dosage:	50 - 100 mg ones daily.
		Side-effects:	Loss of appetite, heartburn and acid regurgitation, nausea and vomiting, abdominal pain, diarrhea, flatulence, headache, dizziness, skin rash, itchy skin.
		Validity:	3 years from the date of production (Voltaren,2care4).
		Storage:	Normal temperature - not above 30°C.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used in case of: 1. allergy to acetylsalicylic acid. 2. asthma patients. 3. patients in blood diluting treatment. 4. patients with symptoms of ulcer.

Grp. 3 Smertestillende midler / Analgetics
(continued)

Stærke smerter	3.9	Lægemiddelstof:	Morphinhydroclorid Δ
		Form:	Tablet 10 mg
		Virkning:	Smertestillende
		Dosering:	10 – 30 mg hver 4 time
		Bivirkninger:	Kvalme, forstoppelse, opkastning, feber, humørsvingninger og lavt blodtryk. Svimmelhed, hallucinationer, nedsat verjtrækning og mavesmerter
		Holdbarhed:	4 år fra produktionsdato (Morfin, EQL Pharma AB)
		Opbevaring:	Normal temperatur. Dog under 25 °C
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i>
Severe pain	3.9	Active Pharmaceutical Ingredients:	Morphine hydroclorid Δ
		Form:	Tablets 10 mg
		Effect:	Pain relieving
		Dosage:	10 – 30 mg every 4 hours
		Side-effects:	Nausea, constipation, vomiting, fever, mood swings and low blood pressure. Dizziness, hallucinations, decreased circulation and abdominal pain
		Validity:	4 years from production date (Morphine, EQL Pharma AB)
		Storage:	Normale temperature – not above 25 °C
		<i>Remarks:</i>	Consult <i>Radio Medical</i>

Grp. 4 Midler mod allergi, shock og søsyge / Agents for Allergy, Shock and Seasickness

Allergisk shock	4.1	Lægemiddelstof:	Adrenalin
		Form:	Injektionsvæske 1 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Øger pulsen, hjertets pumpekraft og blodtrykket. Sammenækning af blodkar. Virker 2 - 5 minutter efter intramuskulær injektion. Omsættes meget hurtigt. Halveringstiden i blodet er 2 - 3 minutter.
		Dosering:	Omgående (voksne): 0,6 mg (godt en halv ampul) dybt intramuskulært. Kan eventuelt gentages, hvis virkningen udebliver.
		Bivirkninger:	Blodtryksstigning, hurtig puls, rysten og svimmelhed.
		Holdbarhed:	2 år fra produktionsdato (Adrenalin®, 2care4),
		Opbevaring:	Læs indlægsseddel
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Er livsfarligt at anvende ved shock forårsaget af blod- eller væsketab.
Allergic shock	4.1	Active Pharmaceutical Ingredients:	Epinephrine
		Form:	Injection liquid, 1 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Increases the pulse, the pumping power of the heart, and the blood pressure. Contraction of blood vessels. Takes effect 2-5 minutes after intramuscular injection. Transformed very quickly. The half-life in the blood is 2-3 minutes.
		Dosage:	Immediately (adults): 0.6 mg (about half an ampoule) given deep intramuscular. May be repeated if the effect is insufficient.
		Side-effects:	High blood pressure, rapid pulse, shivering, and dizziness.
		Validity:	2 years from the date of production (Adrenalin®, 2care4)
		Storage:	Read packet insert
		Remarks:	Consult <i>Radio Medical</i> . Potentially fatal if used to treat shock caused by loss of blood or fluid.

Grp. 4 Midler mod allergi, shock og søsyge / Agents for Allergy, Shock and Seasickness

Allergi Høfeber Nældefeber Allergisk ansigtsødem Allergisk astma	4.2	Lægemiddelstof:	Acrivastin
		Form:	Kapsler à 8 mg.
		Virkning:	Modvirker allergi. Maksimal virkning efter 1½-2 timer. Virkningsvarighed op til 8 timer.
		Dosering:	Voksne: 8 mg efter behov. Maksimalt 3 gange per døgn.
		Bivirkninger:	Døsighed kan forekomme og forstærkes af alkohol og beroligende midler, herunder sovemidler.
		Holdbarhed:	3 år fra produktionsdato (Benadryl®, McNeil ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Lægemidlet er ikke mærket med Δ men enkelte personer sløves alligevel så meget, at det er uforsvarligt med vagttjeneste eller maskinbetjening. Ved lugesymptomer: Se tillige gruppe 15.
Allergy Hay fever Urticaria Allergic face oedema Allergic asthma	4.2	Active Pharmaceutical Ingredients:	Acrivastine
		Form:	Capsule of 8 mg.
		Effect:	Eases allergic symptoms. Maximum effect after 1.5-2 hours. Effective for up to 8 hours.
		Dosage:	Adults: 8 mg when required. Maximum three times daily.
		Side-effects:	Drowsiness may occur and is increased by the use of alcohol and sedatives, including sleeping medicine.
		Validity:	3 years from the date of production (Benadryl®, McNeil Denmark ApS).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . The medicament is not marked with a Δ , but some patients experience a high degree of drowsiness, making bridge duty and operations of machinery unsafe. In case of symptoms of from the lungs: See group 15.
Allergi Allergisk astma Allergisk ansigtsødem	4.3	Lægemiddelstof:	Hydrocortison
		Form:	Injektionssubstans 100 mg i to-delt hætteglas med 2 ml sterilt vand til intramuskulært eller intravenøst brug.
		Virkning:	Modvirker allergi og inflammationer. Virkningen indtræder først 4 - 6 timer efter injektion, hvorfor anden akut behandling skal dække dette tidsrum.
		Dosering:	Voksne: 100 mg intramuskulært eller intravenøst som éngangsdosis.
		Bivirkninger:	Ved længere tids anvendelse: Forstyrrelser i sukker, vand- og saltstofskeft og i kirtlernes funktion. Sløring og/eller aktivering af infektioner. Eufori, sindssyge og søvnløshed. Ondt i maven og mavesår.
		Holdbarhed:	5 år fra produktionsdato (Solu-Cortef®, Pfizer ApS). Brugsfærdig injektionsvæske holdbart i 72 timer, hvis opbevaret i køleskab.
		Opbevaring:	Almindelig temperatur. Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved lugesymptomer: Se tillige gruppe 15.

Grp. 4 Midler mod allergi, shock og søsyge / Agents for Allergy, Shock and Seasickness

Allergy Allergic asthma Allergic face oedema	4.3	Active Pharmaceutical Ingredients:	Hydrocortisonsuccinate
		Form:	Injection substance, 100 mg in capped vial divided into two sections, with 2 ml sterile water for intramuscular or intravenous use.
		Effect:	Eases symptoms of allergy and inflammations. Does not take effect until 4-6 hours after the injection. Other acute treatment to cover this period is needed.
		Dosage:	Adults: 100 mg given intramuscularly or intravenously as a single dose.
		Side-effects:	If used for a extended period: Disturbances in the sugar, water and salt metabolism and in the glandula functions. Masking and/or activation of infections. Euphoria, mental illness and insomnia. Stomacheache and ulcer.
		Validity:	5 years from the date of production (Solu-Cortef®, Pfizer ApS). Injection liquid ready for use, valid for 72 hours, if stored cold.
		Storage:	Normal temperature. To be kept out of daylight. Injection liquid ready for use, to be stored refrigerated.
		Remarks:	Consult <i>Radio Medical</i> . In case of symptoms from lungs See also group 15.
Allergiske reaktioner, hævelser osv. Oral behandling	4.6	Lægemiddelstof:	Prednisolon
		Form:	Tabletter á 25 mg
		Virkning:	Bremser allergiske reaktioner, hævelser osv.
		Dosering:	Efter ordination af <i>Radio Medical</i>
		Bivirkninger:	Forhøjet tryk i øjet, slører infektion og infektion
		Holdbarhed:	2 år fra produktions dato (Prednisolon, Actavis)
		Opbevaring:	Almindelig temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Allergic reactions, swellings etc. Oral treatment preferred	4.6	Active Pharmaceutical ingredients:	Prednisolone
		Form:	Tablets of 25 mg
		Effect:	Brakes allergic reaction, swelling etc.
		Dosage:	After precipstion from <i>Radio Medical</i>
		Side-effects:	Glaucoma, blurs infections and infekctions
		Validity:	2 years from productions date (Prednisolon, Actavis)
		Storage:	Normal temperature
		Remarks:	Consult <i>Radio Medical</i>

Grp. 4 Midler mod allergi, shock og søsyge / Agents for Allergy, Shock and Seasickness

Søsyge	4.7	Lægemiddelstof:	Cyclizine Δ
		Form:	Table á 50 mg
		Virkning:	Modvirker søsyge
		Dosering:	50 mg, 1-2 timer før ønsket effekt. Højest 3 doser pr. døgn
		Bivirkninger:	Sløret syn, Døshed
		Holdbarhed:	3 år fra produktiondato (Marzine, Viatris ApS)
		Opbevaring:	Normal temperatur
		Anmærkninger:	Søg samråd med Radio Medical
Seasickness	4.7	Active Pharmaceutical ingredients:	Cyclizine Δ
		Form:	Tablets of 50 mg
		Effect:	Prevent seasickness
		Dosage:	50 mg 1-2 hours before effect. Maximum 3 dosage pr. 24 hours
		Side-effects:	Blurred vision, drowsiness
		Validity:	3 years from production date (Marzine, Viatris ApS)
		Storage:	Normale temperature
		Remarks:	Consult Radio Medical

Grp. 5 Antidoter / Antidotes

Modgift ved overdosering af morfin MFAG tabel 4, 13 Førstehjælp: Ved udsættende vejtrækning understøttes manuelt med ilt på ballon og maske 15 l ilt/minut.	5.1	Lægemiddelstof:	Naloxon
		Form:	Injektionsvæske 0,4 mg/ml, ampuller á 1 ml til intramuskulær injektion.
		Virkning:	Naloxon ophæver morfins (bi)virkning. Virker 2 - 3 minutter efter intramuskulær injektion. Virkningsvarighed: Op til 4 timer.
		Dosering:	1. Naloxon voksne: Der gives omgående 0,8 mg (2 ampuller) intramuskulært. 2. Hvis virkningen udebliver kan behandlingen gentages efter 5 minutter.
		Bivirkninger:	Rysten, utilpashed og opkastninger (abstinenssymptomer).
		Holdbarhed:	3 år fra produktionsdato (Naloxon, B. Braun).
		Opbevaring:	Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Antidote to be administered in case of an overdose of morphine MFAG table 4, 13 First Aid: If the patient stops breathing, assist breathing manually with oxygen 15l/minute on a mask.	5.1	Active Pharmaceutical Ingredients:	Naloxone
		Form:	Injection liquid, 0.4 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Naloxone counteracts the (side)-effects of morphine. Takes effect 2-3 minutes after intramuscular injection. Effective for up to 4 hours.
		Dosage:	1. Naloxone adults: 0.8 mg (2 ampoules) is given immediately intramuscular. 2. In case of no effect, the treatment may be repeated after 5 minutes.
		Side-effects:	Shivering, feeling unwell, and vomiting.
		Validity:	3 years from the date of production (Naloxon B. Braun).
		Storage:	To be protected from daylight.
		Remarks:	Consult <i>Radio Medical</i> .

Insulintilfælde Insulinshock Lavt blodsukker	5.2	Lægemiddelstof:	Glucagon
		Form:	Ready-set bestående af: Injektionssubstans Glucagon 1 mg i hætteglas og sprøjte med 1 ml sterilt vand til intramuskulært brug.
		Virkning:	Frigør sukkerreserver fra leveren. Virker normalt inden 5 - 10 minutter. Virkningsvarighed: 10 - 40 minutter.
		Dosering:	Voksne: 1 mg intramuskulært. Kan evt. gentages efter 10 - 15 minutter. Når patienten er kommet til bevidsthed gives brød og sød saft.
		Bivirkninger:	Kvalme og opkastninger kan forekomme. Kan udløse hjertekrampe hos hjertesygge. Hjertekrampe behandles med glycerylnitrat spray (9.1).
		Holdbarhed:	Ved almindelig temperatur: 18 måneder fra produktionsdato (GlucaGen® HypoKit, Novo Nordisk). Ved 2 - 8° C: 36 måneder fra produktionsdato. Den blandede opløsning skal anvendes umiddelbart.
		Opbevaring:	Almindelig temperatur eller 2 - 8° C. (GlucaGen® HypoKit, Novo Nordisk).
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Dersom den blandede opløsning er uklar eller med krystaller, må blandingen ikke anvendes.
Hypoglycaemic coma Insulin shock Low blood sugar	5.2	Active Pharmaceutical Ingredients:	Glucagon
		Form:	Ready-set consisting of: Injection substance Glucagon, 1 mg in a capped vial and a syringe with 1 ml of sterile water for intramuscular use.
		Effect:	Releases sugar reserves from the liver. Normally, takes effect within 5-10 minutes. Effective for 10-40 minutes.
		Dosage:	Adults: 1 mg intramuscularly. May be repeated after 10-15 minutes. When the patient regains consciousness, he is given some bread and sweet juice.
		Side-effects:	Nausea and vomiting may occur. May cause angina pectoris in people with heart diseases. Angina pectoris is treated with glyceryl trinitrate spray (9.1).
		Validity:	At room temperature: 18 months from the date of production (GlucaGen® HypoKit, Novo Nordisk). At 2-8°C: 36 months from the date of production. The mixed dissolution must be used immediately.
		Storage:	Normal temperature or 2-8° C. (GlucaGen® HypoKit, Novo Nordisk).
		Remarks:	Consult <i>Radio Medical</i> . If the mixed dissolution is cloudy or contains crystals, it must not be used.

MFAG tabel 17 Forgiftninger med insektgifte, der indeholder organophosphater eller carbamater.	5.3	Lægemiddelstof:	Atropin
		Form:	Injektionsvæske 1 mg/ml, ampuller à 1 ml til intravenøs injektion.
		Virkning:	Livreddende ved forgiftninger af visse insektgifte. Virkningen afhænger af gifttype og mængde. Se MFAG tabel 17.
		Dosering:	Dosis afhænger af forgiftningen. Atropin indgives langsomt intravenøst. Der startes med 2 mg (2 ampuller) og fortsættes med yderligere 2 mg efter hver 5 - 10 minutters mellemrum indtil pupillerne begynder at udvide sig. Giftig dosis er 10 - 20 mg til voksne. Dødelig dosis er ca. 200 mg til voksne.
		Bivirkninger:	Pupiludvidelse, lysskyhed, tørre slimhinder, hurtig puls, vandladningsbesvær og forstoppelse.
		Holdbarhed:	Plastampuller: 2 år fra produktionsdato. (Atropin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
MFAG table 17 Poisoning with insecticides containing organophosphates or carbamates.	5.3	Active Pharmaceutical Ingredients:	Atropin
		Form:	Injection liquid, 1 mg/ml, ampoules of 1 ml for intravenous injection.
		Effect:	Life-saving in connection with poisonings caused by certain insecticides. The effect depends on the type and amount of poison. See MFAG table 17.
		Dosage:	The dose depends on the poisoning. Atropin is given slowly intravenously. At first, 2 mg (2 ampoules) is given, followed by another 2 mg after every 5-10 minutes until the pupils start dialating. Poisonous dose is 10-20 mg for adults. Lethal dose is about 200 mg for adults.
		Side-effects:	Dilation of the pupils, photophobia, dry mucous membranes, rapid pulse, dysuria and constipation.
		Validity:	Plastic ampoules: 2 years from the date of production (Atropin®, Nycomed Danmark)
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
MFAG tabel 8, 16 Ætsninger med fluorholdige salte og syrer. Første hjælp: Huden skylles med rigelig vand i mindst 10 minutter. Forurenet tøj fjernes. Huden vaskes med vand og sæbe og skylles.	5.4	Lægemiddelstof:	Calciumgluconat 2%
		Form:	Tuber à 25 g, indeholdende 2 g calciumgluconat per 100 g. Fremstilles af apoteket.
		Virkning:	Calcium neutraliserer i et vist omfang ætsningsskaderne ved fluoridætsninger.
		Dosering:	Huden indsmøres i calciumgluconat gel. Se vejledning i MFAG tabel 8 og 16.
		Bivirkninger:	Ingen.
		Holdbarhed:	2 år fra produktionsdato.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .

MFAG table 8, 16 Caustic burns caused by salts and acids containing fluorine.	5.4	Active Pharmaceutical Ingredients:	Calciumgluconate 2%
		Form:	Tubes of 25 g, containing 2 g of calciumgluconate per 100 g. Made by the individual pharmacy.
		Effect:	Calcium neutralizes caustic burns caused by fluorine to some extent.
		Dosage:	Calciumgluconate gel is applied to the skin. See guidelines in MFAG table 8 og 16.
		Side-effects:	None.
		Validity:	2 years from the date of production.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
MFAG tabel 16 Ætsninger med fluorholdige salte og syrer, der omfatter mere end 1% af hudens overflade.	5.5	Lægemiddelstof:	Calciumlactogluconat og calciumcarbonat.
		Form:	Brusetabletter à 500 mg.
		Virkning:	Neutraliserer i et vist omfang skader efter optagelse af fluorsalte og fluorid forbindelser gennem huden.
		Dosering:	Ordination af <i>Radio Medical</i> .
		Bivirkninger:	Normalt ingen bivirkninger.
		Holdbarhed:	3 år fra produktionsdato (Calcium-Sandoz®, Novartis).
		Opbevaring:	Er fugtfølsomt. Opbevares tørt i tæt emballage ved almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
MFAG table 16 Caustic burns caused by salts and acids containing fluorine and covering more than 1% of the surface of the skin.	5.5	Active Pharmaceutical Ingredients:	Calciumlactogluconate and calciumcarbonate
		Form:	Effervescent tablets of 500 mg.
		Effect:	Neutralizes injuries caused by the absorption of fluoride compounds through the skin to some extent.
		Dosage:	Contact <i>Radio Medical</i> .
		Side-effects:	No side effects.
		Validity:	3 years from the date of production (Calcium-Sandoz, Orifarm).
		Storage:	Sensitive to humidity. To be stored in a dry place in a sealed packaging at room temperature.
		Remarks:	Consult <i>Radio Medical</i> .

MFAG tabel 10 Forgiftninger og ætsninger i mave og tarm.	5.6	Lægemiddelstof:	Aktivt kul.
		Form:	50 g aktivt kul i granulatform til opblanding i 400 ml vand.
		Virkning:	Opsuger gift og lægemidler fra mavetarmkanalen og binder stofferne til kulpartiklerne.
		Dosering:	50 - 100 g kul (1 - 2 flasker) opslemmet i vand indtages så hurtigt som muligt. Ved alvorlige forgiftninger fortsættes med 20 g (160 ml) hver 6. time. Se også MFAG, Appendix 14.
		Bivirkninger:	Farver afføringen sort. Kan give forstoppelse. Nedsætter virkningen af lægemidler, der er indtaget via munden.
		Holdbarhed:	5 år fra produktionsdato (Carbomix Norit®). Efter opblanding i vand: Maks. 3 dage i køleskab.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Anvendes ikke til bevidstløse. Må ikke anvendes ved indtagelse af stærke syrer/baser eller organiske opløsningsmidler (petroleums-lignende).
MFAG table 10 Poisoning and caustic burns in stomach and intestines.	5.6	Active Pharmaceutical Ingredients:	Active carbon
		Form:	50 g of active carbon in the form of granulate for dilution in 400 ml of water.
		Effect:	Absorbs poison and medicaments from the gastrointestinal tract and binds the substances to the carbon particles.
		Dosage:	50-100 g carbon (1-2 bottles) suspended in water is taken as quickly as possible. In case of serious poisonings, continue with 20 g (160 ml) every sixth hour. See also MFAG, Appendix 14.
		Side-effects:	Black faeces. May cause constipation. Reduces the effect of medicaments taken through the mouth.
		Validity:	5 years from the date of production (Carbomix Norit®). After having been mixed with water, maximum 3 days in the fridge.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Do not use on unconscious patients. Not to be used in connection with strong acids/bases or organic solvents (kerosene related).

MFAG tabel 19 Forgiftninger med methanol og ethylene glycol	5.7	Lægemiddelstof:	Ethyl alkohol 99,5% (alternativt 99,9%).
		Form:	Væske
		Virkning:	Modgift mod methanol og ethylenglycols virkning.
		Dosering:	25 ml opblandes i 250 – 300 ml vand eller saft. Dosis gentages hver 3 time indtil evakuering. Se MFAG tabel 19.
		Bivirkninger:	Kvalme, hovedpine.
		Holdbarhed:	2 år fra produktionsdato.
		Opbevaring:	Almindelig stuetemperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
MFAG table 19 Poisoning caused by methanole and ethylene glycole	5.7	Active Pharmaceutical Ingredients:	Ethyl alcohol 99,5% (alternative 99,9%)
		Form:	Liquid
		Effect:	Antidote against the effect of methanole and ethylene glycole.
		Dosage:	25 ml is mixed with 250-300 ml water or juice. Repeat every 3 hours until evacuation. See MFAG table 19.
		Side-effects:	Nausea, vomiting and headache.
		Validity:	2 years from the date of production.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
MFAG tabel 14 Forgiftning med medikamenter (acetylsalicylsyre) og pesticider med blodfortyndende virkning (coumarinderivater, warfarrin, bromadiolon m.fl.)	5.8	Lægemiddelstof:	Phytomenadion (K-vitamin)
		Form:	Injektionsvæske 10 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Modvirker gifterter, der medfører forstyrrelser i blodets evne til at størkne. Maksimal virkning 2 timer efter injektion.
		Dosering:	Se MFAG tabel 14. Sædvanlig dosis: 10 - 20 mg dagligt.
		Bivirkninger:	Udvidelse af blodkarrene, blodtryksfald, åndenød, kollaps.
		Holdbarhed:	3 år fra produktionsdato (Konakion® Novum, Roche).
		Opbevaring:	Almindelig temperatur, beskyttet mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Stoffet må ikke anvendes, hvis væsken er uklar eller har delt sig i faser.
MFAG table 14 Poisoning caused by medicaments (acetylsalicylic acid) and pesticides with a blood- thinning effect (coumarine derivates, warfarin, bromadiolone, etc.)	5.8	Active Pharmaceutical Ingredients:	Phytomenadione (vitamin K)
		Form:	Injection liquid, 10 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Counteracts types of poison causing disturbances in the blood's ability to coagulate. Maximum effect 2 hours after injection.
		Dosage:	See MFAG table 14. Ordinary dose: 10-20 mg daily.
		Side-effects:	Enlargement of the blood vessels, falling blood pressure, difficulty in breathing, collapse.
		Validity:	3 years from the date of production (Konakion®, Novum, Roche).
		Storage:	Normal temperature, protected from daylight.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used if the liquid is cloudy or has divided itself into phases.

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion

Angst og urotilstande Alkoholabstinenser Forebyggelse af kramper Ledskred og brud 1. valgspræparat ved akut sindssyge (alternativt anvendes klysma (6.3))	6.2	Lægemiddelstof:	Diazepam ^Δ
		Form:	Tabletter à 5 mg.
		Virkning:	Stærkt sløvende og muskelafslappende. Dæmper angst, uro og rastløshed. Virker efter 30 minutter. Udskilles meget langsomt (Halveringstid: 20 - 50 timer). Forstærker virkningen af morfin, haloperidol, alkohol og andre sløvende medikamenter.
		Dosering:	5 mg x 3 dagligt. Akut sindssyge: Dosis efter aftale med <i>Radio Medical</i> . Se også 6.4.
		Bivirkninger:	Vanedannende. Døsighed, svimmelhed, forvirring, hukommelsessvigt og balanceforstyrrelser.
		Holdbarhed:	5 år fra produktionsdato (Diazepam, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Anxiety and states of restlessness Alcoholic withdrawal symptoms Prevention of convulsions Dislocations and fractures First choice of medicament in case of acute mental disease (alternatively, Enema (6.3) are used.)	6.2	Active Pharmaceutical Ingredients:	Diazepam ^Δ
		Form:	Tablets of 5 mg.
		Effect:	Strongly sedative and muscle-relaxant. Decreases anxiety and restlessness. Takes effect after 30 minutes. Is excreted very slowly (halving time: 20-50 hours). Increases the effect of morphine, haloperidole, alcohol and other sedatives.
		Dosage:	5 mg three times daily Acute mental disease: Dose after consultation with <i>Radio Medical</i> . See also 6.4.
		Side-effects:	Addictive. Drowsiness, dizziness, confusion, lapse of memory, and balance disturbances.
		Validity:	5 years from the date of production (Diazepam, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Universelle kramper Kramper fra forgiftninger Abstinenskramper Epilepsi Akut sindssyge MFAG table 4, 5, 6	6.3	Lægemiddelstof:	Diazepam ^Δ
		Form:	Klysma, 4 mg/ml, klysma til anvendelse i endetarmen
		Virkning:	Stærkt sløvende og muskelafslappende. Dæmper angst, uro og rastløshed. Virker 5 - 10 minutter efter rektal administration. Udskilles meget langsomt (Halveringstid: 20 - 50 timer). Forstærker virkningen af morfin, haloperidol, alkohol og andre sløvende medikamenter.
		Dosering:	Efter aftale med <i>Radio Medical</i> - 1 klysma tømmes ind i endetarmen.
		Bivirkninger:	Døsighed, svimmelhed, forvirring, hukommelsessvigt og balanceforstyrrelser.
		Holdbarhed:	2,5 år fra produktionsdato (Stesolid®, Actavis A/S). Dog maksimalt 1 år efter anbrud af aluminiumsposen.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion (continued)

Universal convulsions Convulsions caused by poisoning Withdrawal convulsions Epilepsy Acute mental illness MFAG tabel 4, 5, 6	6.3	Active Pharmaceutical Ingredients:	Diazepam Δ
		Form:	Enema, 4 mg /ml, enema for insertion into the rectum.
		Effect:	Strong sedative and muscle-relaxant. Decreases anxiety and restlessness. Takes effect 5-10 minutes after rectal administration. Is excreted very slowly (half-life: 20-50 hours). Increases the effect of morphine, haloperidole, alcohol and other sedatives.
		Dosage:	After consult of <i>Radio Medical</i> - 1 enema is emptied into the rectum.
		Side-effects:	Drowsiness, dizziness, confusion, lapse of memory, and balance disturbances.
		Validity:	2,5 years from the date of production (Stesolid®, Actavis A/S). However, maximum 1 year after having opened the Aluminiumbag.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
2. valgspræparat ved akut sindssyge præget af mani, vildskab, psykisk og Motorisk uro, angst, vrangforestillinger og hallucinationer. Kvalme og opkastninger	6.4	Lægemiddelstof:	Haloperidol Δ
		Form:	Injektionsvæske 5 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Afdæmper symptomerne ved sindssygen. Dæmper kvalme og opkastninger.
		Dosering:	Akut sindssyge: 5 - 10 mg intramuskulært. Kan eventuelt gentages 3 eller flere gange dagligt. Behandlingen kan eventuelt forstærkes med diazepam (6.2/ 6.3).
		Bivirkninger:	Den sløvende effekt forstærkes af alkohol, morfin og andre sløvende midler.
		Holdbarhed:	5 år fra produktionsdato (Serenase®, Janssen-Cilag).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Second choice of medicament in case of acute mental illness characterized by mania, wildness, mental and motoric restlessness, anxiety, delusions, and hallucinations. Nausea and vomiting	6.4	Active Pharmaceutical Ingredients:	Haloperidole Δ
		Form:	Injection liquid 5 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Restricts the symptoms of mental disease. Reduces nausea and vomiting.
		Dosage:	Acute mental disease: 5-10 mg intramuscularly. May be repeated three times daily or more. The treatment may be enhanced by the use of diazepam (6.2/6.3).
		Side-effects:	The sedative effect is increased by alcohol, morphine, and other sedatives.
		Validity:	5 years from the date of production (Serenase®, Janssen-Cilag).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion (continued)

Søvnløshed Indsovningsbesvær	6.5	Lægemiddelstof:	Zolpidem Δ
		Form:	Tabletter à 10 mg.
		Virkning:	Stærkt sløvende. Virker hurtigt efter indtagelse. Virkningen varer op til 6 timer. Virkningen forstærkes af andre sløvende stoffer, herunder alkohol.
		Dosering:	Voksne. 10 mg 15-30 min. før sengetid.
		Bivirkninger:	Kvalme, opkastning, diarré. Hovedpine, koncentrationsbesvær, konfusion, hukommelsesbesvær, svimmelhed, døsigheid, hudreaktioner, dobbeltsyn.
		Holdbarhed:	5 år fra produktionsdato (Zolpidem Hexal A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg råd med <i>Radio Medical</i> .
Insomnia Difficulty falling asleep	6.5	Active Pharmaceutical	Zolpidem Δ
		Form:	Tablets of 10 mg.
		Effect:	Strong sedative, rapid effect after ingestion. Effective up till 6 hours. The effect is increased by the use of other sedatives, including alcohol.
		Dosage:	Adults. 10 mg 15-30 minutes. before bedtime.
		Side-effects:	Nausea, vomiting, diarrhea. Headache, difficulty concentrating, confusion, memory loss, dizziness, drowsiness, skin reactions, double vision.
		Validity:	5 years from the date of production (Zolpidem Hexal A/S).
		Storage:	Normal temperature.

Grp. 7 Midler mod infektioner / Infections			
Halsbetændelse Mundhuleinfektioner Bihulebetændelse Mellemørebetændelse Lungebetændelse	7.1	Lægemiddelstof:	Phenoxymethylpencillin
		Form:	Tabletter à 660 - 800 mg (ca.1 mill. IE).
		Virkning:	Dræber de hyppigst forekommende bakterier, der forårsager infektioner i luftvejene.
		Dosering:	400–800 mg × 2 – 4 dagligt.
		Bivirkninger:	Kvalme, diarré og hududslæt. Sjældent ses overfølsomhedsshock.
		Holdbarhed:	3 år fra produktionsdato (Primcillin®, MEDA).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg råd med <i>Radio Medical</i> . Overfølsomhedsshock optræder meget sjældent, og som regel senest 30 minutter efter indgift. Behandling: Adrenalin 0.6 mg intramuskulært (4.1).
Inflammation of the throat Infections of the oral cavity Sinusitis Inflammation of the middle ear Pneumonia	7.1	Active Pharmaceutical	Phenoxymethylpenicillin
		Form:	Tablets of 660 - 800 mg (approx. 1 MIU).
		Effect:	Kills the most common bacteria causing infections of the respiratory tract.
		Dosage:	400-800 mg × 2-4 times daily.
		Side-effects:	Vomiting, diarrhoea and rash. Rarely allergic shock.
		Validity:	3 years from the date of production (Primcillin®, MEDA).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Allergic shock is very rare and usually occurs within 30 minutes after administration. Treatment: Adrenaline 0.6mg intramuscular.(4.1).
MFAG tabel 9 Lungebetændelse efter inhalation af giftige kemiske stoffer	7.2	Lægemiddelstof:	Amoxicillin
		Form:	Kapsler eller tabletter à 500 mg
		Virkning:	Modvirker inflammation
		Dosering:	375-750 mg 2 gange daglig
		Bivirkninger:	Feber, kvalme, opkastning, sure opstød, diarré, hudkløe og hududslæt
		Holdbarhed:	4 år fra produktionsdato (Imadrax, Sandoz A/S)
		Opbevaring:	Almindelig temperatur
		Anmærkninger:	Søg råd med <i>Radio Medical</i> Overfølsomhedsshock optræder meget sjældent og som regel senest 30 minutter efter indgift. Behandling: Adrenalin 0,6 mg intramuskulært (4.1)
MFAG table 9 Pneumonia after inhalation of poisonous chemical substances	7.2	Active Pharmaceutical Ingredients:	Amoxicilline
		Form:	Capsules or tablets of 500 mg
		Effect:	Prevent inflammation
		Dosage:	375-750 mg twice daily
		Side-effects:	Fever, nausea, vomiting, heartburn, diarrhoea, rash and urticaria, Rarely allergic shock
		Validity:	4 years from the date of production (Imadrax, Sandoz A/S)
		Storage:	Normal temperature
		Remarks:	Consult <i>Radio Medical</i> . Allergic shock is very rare and usually it occurs within 30 minutes after administration. Treatment: Adrenaline 0.6 mg intramuscular (4.1)

Grp. 7 Midler mod infektioner / Infections				(continued)
Halsbyld Tandbyld Knogleinfektion Underlivsinfektioner Tarminfektioner Blindtarmsbetændelse (tjek urinstiks G.1)	7.6	Lægemiddelstof:	Metronidazol	
		Form:	Tabletter à 500 mg.	
		Virkning:	Bakteriedræbende.	
		Dosering:	500 mg x 3 dagligt.	
		Bivirkninger:	Mavekneb, smagsforstyrrelser, hovedpine, hududslet og svimmelhed. Urinen kan blive farvet mørk. Kan virke som antabus ved samtidig indtagelse af alkohol.	
		Holdbarhed:	5 år fra produktionsdato (Metronidazol, Actavis).	
		Opbevaring:	Almindelig temperatur.	
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes til gravide. Anvendes ofte i kombination med andre antibiotika, f.eks. Ciprofloxacin (7.8) eller Ceftriaxon (7.9) Må ikke anvendes sammen med 15.1 Dextromethorphan.	
Boil in the throat Tooth abscess Bone infection Infection of female reproductions organs Abdominal infections Intestinal infections Appendicitis (Check dipstick G.1)	7.6	Active Pharmaceutical Ingredients:	Metronidazole	
		Form:	Tablet of 500 mg.	
		Effect:	Kills bacteria.	
		Dosage:	500 mg three times daily.	
		Side-effects:	Bellyache, altered sense of taste, headache, rash, and dizziness. The urine may be coloured black. May cause a reaction similar to that caused by Antabuse in connection with alcohol.	
		Validity:	5 years from the date of production (Metronidazol, Actavis).	
		Storage:	Normal temperature.	
		Remarks:	Consult <i>Radio Medical</i> .	
Halsbyld Tandbyld Knogleinfektion Underlivsinfektioner Tarminfektioner Blindtarmsbetændelse MFAG tabel 10	7.7	Lægemiddelstof:	Metronidazol	
		Form:	Stikpille à 500 mg.	
		Virkning:	Bakteriedræbende.	
		Dosering:	500 mg x 3 daglig.	
		Bivirkninger:	Kvalme, opkastning, mavesmerter, diarré, forstoppelse, og smagsforstyrrelse. Mørkfarvet urin. Kan virke som antabus ved samtidig indtagelse af alkohol.	
		Holdbarhed:	3 år fra produktionsdato (Flagyl® , Sanofi-aventis)	
		Opbevaring:	Almindelig temperatur.	
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan om nødvendigt anvendes, til gravide. Anvendes ofte i kombination med andre antibiotika, fx Ciprofloxacin (7.8) eller Ceftriaxon (7.9). Må ikke anvendes sammen med 15.1 Dexofan.	

Grp. 7 Midler mod infektioner / Infections				(continued)
Boil in the throat Tooth abscess Bone infection Infection of female reproductions organs Abdominal infections Intestinal infections Appendicitis (Check dipstick G.1) MFAG table 10	7.7	Active Pharmaceutical Ingredients:	Metronidazol	
		Form:	Suppository of 500 mg.	
		Effect:	Kills bacteria.	
		Dosage:	500 mg three times daily.	
		Side-effects:	Nausea vomiting, abdominal pain, diarrhea, constipation, and taste disturbance. Dark colored urine. May cause a reaction similar to Antabuse in connection with alcohol.	
		Validity:	3 years from productions date (Flagyl®, Sanofi-aventis	
		Storage:	Normal temperature.	
		Remarks:	Consult <i>Radio Medical</i> . May be used during pregnancy, if nessasary. Often used in combination with other antibiotics e.g. Ciprofloxacin (7.8) or Ceftriaxon (7.9). May not be used with 15.1 Dextromethorphan.	
Tandbyld Halsbyld Svær bronkitis Lungepest Akut diarré Kolera Dysenteri Blindtarmsbetændelse Gonoré Urinrørsbetændelse Nyrebækkenbetændelse (tjek urinstiks G.1)	7.8	Lægemiddelstof:	Ciprofloxacin	
		Form:	Tabletter à 250 mg.	
		Virkning:	Dræber bakterier.	
		Dosering:	250 - 750 mg x 2 dagligt.	
		Bivirkninger:	Mavekneb, kvalme, opkastninger, madlede, hovedpine, svimmelhed, træthed og kramper, muskel-, led- og seneømhed, allergi, overfølsomhedsreaktion ved samtidig	
		Holdbarhed:	5 år fra produktionsdato (Cifin, PharmaCoDane ApS).	
		Opbevaring:	Almindelig temperatur.	
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Gives ved mange infektioner i kombination med metronidazol (7.6 eller 7.7). Må ikke gives til gravide eller ammende.	
Boil in the throat Tooth abscess Severe bronchitis Pneumonic plague Acute diarrhoea Cholera Dysentery Appendicitis Gonorrhoea Urethritis Pyelitis (Check dipstick G.1	7.8	Active Pharmaceutical Ingredients:	Ciprofloxacin	
		Form:	Tablets of 250 mg.	
		Effect:	Kills bacteria.	
		Dosage:	250-750 mg twice dailyThe tablets are given 1 hour before a	
		Side-effects:	Bellyache, nausea, vomiting, loss of appetite, headache, dizziness, fatigue, and convulsions. Soreness in the muscles, joints and tendons. Allergy. Allergic reaction in case of simultaneous exposure	
		Validity:	5 years from the date of production (Cifin, PharmaCoDane Aps).	
		Storage:	Normal temperature.	
		Remarks:	Consult <i>Radio Medical</i> . Is given in combination with metronidazole (7.6 or 7.7) in connection with many infections. Should not be given to pregnant or breast-feeding patients.	

Grp. 7 Midler mod infektioner / Infections			(continued)
Blodforgiftning Sepsis Meningitis Lungebetændelse Blindtarmsbetændelse Nyrebækkenbetændelse Store sårskader Åbne knoglebrud MFAG tabel 10 (Ceftriaxon erstatter Cefuroxime)	7.9	Lægemiddelstof:	Ceftriaxon
		Form:	Injektionssubstans, 1 g tørstof i hætteglas. Kan gives intramuskulært - Kontakt <i>Radio Medical</i> .
		Virkning:	Bakteriedræbende.
		Dosering:	1 - 2 g intravenøst i døgnet i én dosis.
		Bivirkninger:	Allergiske reaktioner: Hududslet, nældefeber, ledsmerter, overfølsomhedsshock.
		Holdbarhed:	Injektionssubstans: 3 år fra produktionsdato (Rocephalin®, Roche). Brugsfærdig opløsning: 24 timer i køleskab.
		Opbevaring:	Må ikke opbevares over 30°C.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Gives ved mange infektioner i kombination med metronidazol (7.6 eller 7.7). Er stærkt vævsirriterende ved intramuskulær injektion.
Sepsis Blood poisoning Meningitis Pneumonia Appendicitis Pyelitis Major wounds Open fractures MFAG table 10 (Ceftriaxone replaces Cefuroxime)	7.9	Active Pharmaceutical Ingredients:	Ceftriaxone
		Form:	Injection substance g of dry substance in a capped vial. Must be mixed with 10 ml of sterile water (16.7) Can be given intramuscular - Contact <i>Radio Medical</i>
		Effect:	Kills bacteria.
		Dosage:	1-2 g intravenously dailyas a single dose.
		Side-effects:	Allergic reactions: Rash, urticaria, arthralgia, allergic shock.
		Validity:	Injection substance: 3 years from the date of production (Rocephaline®, Roche). Dissolution ready for use: 24 hours if refrigerated
		Storage:	Not to be stored at more than 30°C.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Often used in combination with metronidazole (7.6 or 7.7) in connection with many infections. In case of intramuscular injection, strong irritation of tissue.
Stivkrampevaccine Gives ved ulykker med kvæstelser og sår, med mindre den skadede med sikkerhed er dækket med stivkrampe-vaccine. Vaccinen suppleres med tetanus immunglobulin (se 7.11). Vaccination skal ske i behandlingsrum med adgang til adrenalin (4.1) og genoplivningsudstyr (A.1).	7.10	Lægemiddelstof:	Tetanus holdig vaccine (tetanustoxoid)
		Form:	Injektionsvæske
		Virkning:	Aktiverer patientens immunsystem.
		Dosering:	Følg ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Lokal rødme, hævelse og ømhed. Hovedpine, utilpashed og feber ses undertiden og fortager sig efter et døgn
		Holdbarhed:	2 år fra produktionsdato (Tetanusvaccine SSI).
		Opbevaring:	Opbevares køligt mellem 2 - 8° C. Frossen vaccine må ikke anvendes
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Overfølsomhedsshock optræder sjældent. Behandling: Adrenalin 0,6 mg intramuskulært 4.1).

Grp. 7 Midler mod infektioner / Infections				(continued)
Tetanus vaccine To be administered in case of accidents causing injury and wounds unless the wounded person is for certain covered by a tetanus vaccine. The vaccine is to be supplemented by tetanus immune globulin (see 7.11). The vaccination must be given in a room with access to Epinephrine (4.1) and Resuscitation equipment (A.1).	7.10	Active Pharmaceutical Ingredients:	Vaccine containing Tetanus (tetanustoxoid)	
		Form:	Injection liquid	
		Effect:	Activates the patient's immune system.	
		Dosage:	Follow prescription from <i>Radio Medical</i> .	
		Side-effects:	Local blushing, swelling, and soreness.	
		Validity:	2 years from the date of production (Tetanus vaccine SSI).	
		Storage:	To be stored cool at between 2-8°C. Frozen vaccine must not	
		Remarks:	Consult <i>Radio Medical</i> .	
Stivkrampemodgift Gives ved ulykker med kvæstelser og sår, med mindre den skadede med sikkerhed er dækket med stivkrampe-vaccine. Tetanus immunglobulin suppleres med vaccine (se 7.10). Behandlingen skal ske i behandlingsrum med adgang til 4.1 Adrenalin og A.1 Genoplivnings- udstyr.	7.11	Lægemiddelstof:	Humant tetanus immunglobulin	
		Form:	Injektionsvæske 250 I.E. Ampul à 1 ml til intramuskulær injektion.	
		Virkning:	Bekæmper giftstoffer produceret af stivkrampebakterien.	
		Dosering:	1 injektion à 250 I.E. (1 ml) intramuskulært. Suppleres med	
		Bivirkninger:	Lokal rødme, hævelse, ømhed og feber ses undertiden. Allergisk shock og andre allergiske reaktioner forekommer, men sjældent.	
		Holdbarhed:	3 år fra produktionsdato (Tetagam® P/Tetanogamma® P,	
		Opbevaring:	Opbevares køligt mellem 2 - 8° C. Frossen immunglobulin må ikke anvendes. Må ikke anvendes, hvis væsken fremstår uklar eller klumpet.	
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Overfølsomhedsshock optræder sjældent. Behandling: Adrenalin 0,6 mg intramuskulært (4.1).	
Tetanus antidote To be administered in case of accidents causing injury and wounds unless the wounded person is for certain covered by a tetanus vaccine. Tetanus immune globulin is to be supplemented by a vaccine (see 7.10). The treatment must be given in a room with access to 4.1 Adrenaline and A.1 Resuscitation equipment.	7.11	Active Pharmaceutical Ingredients:	Human tetanus immunoglobulin	
		Form:	Injection liquid 250 IU. Ampoule of 1 ml for intramuscular injection.	
		Effect:	Combats poisonous substances produced by the tetanus	
		Dosage:	1 injection of 250 IU (1 ml) given intramuscularly. To be supplemented by the tetanus vaccine (see 7.10).	
		Side-effects:	Local blushing, swollenness, soreness, and fever are sometimes seen.	
		Validity:	3 years from the date of production (Tetagam® P/ Tetanogamma® P, CSL Behring).	
		Storage:	To be stored cool at between 2-8°C. Frozen immune globulin must not be used. Not to be used if the liquid is muddy or lumpy.	
		Remarks:	Consult <i>Radio Medical</i> .	

Grp. 7 Midler mod infektioner / Infections			(continued)
Infektioner forårsaget af stafylokokker. Knogleinfektion	7.12	Lægemiddelstof:	Flucloxacillin
		Form:	Tabletter à 500 mg.
		Virkning:	Bakteriedræbende
		Dosering:	750 - 1000 mg 3-4 gange dagligt.
		Bivirkninger:	Kvalme, opkastning, diarré, hududslæt.
		Holdbarhed:	2 år fra produktionsdato. Åbne blisterpakninger 6 måneder (Heracillin, Meda AS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Infections caused by staphylococci. Infections in the bones	7.12	Active Pharmaceutical Ingredients:	Flucloxacillin
		Form:	Tablets of 500 mg.
		Effect:	Kills bacteria.
		Dosage:	750-1000 mg 3-4 times daily.
		Side-effects:	Nausea, vomiting, diarrhea, skin rash.
		Validity:	2 years from the date of production. 6 months for open blister pack (Heracillin, Meda AS).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Halsbetændelse Mundhuleinfektioner Bihulebetændelse Mellemørebetændelse Lungebetændelse Legionella Chlamydia infektioner Mavesårsinfektioner Alternativ til penicillin ved allergi MFAG tabel 9	7.15	Lægemiddelstof:	Azithromycin
		Form:	Tabletter á 500 mg
		Virkning:	Standser bakteriernes vækst
		Dosering:	Voksne: 500 mg x 1-2 dagligt
		Bivirkninger:	Mavesmerter, kvalme, nedsat appetit, kraftsløshed, svimmelhed, smagsforstyrrelser, hovedpine, hudkløe og hududslæt
		Holdbarhed:	5 år fra produktionsdato (Azithromycin, KRKA)
		Opbevaring:	Almindelig temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Inflammation of the throat Infections of the oral cavity Sinusitis Inflammation of the middle ear Pneumonia Legionella Chlamydia infections Infections in connection with ulcer Alternative to penicillin in case of allergy MFAG table 9	7.15	Active Pharmaceutical ingredients:	Azithromycin
		Form:	Tablets of 500 mg
		Effect:	Stops bacterial growth
		Dosage:	Adults: 500 mg x 1-2 daily
		Side-effects:	Abdominal pain, nausea, decreased appetite, weakness, dizziness, taste disturbances, headache, itchy skin and skin rash
		Validity:	5 years from date of production (Azithromycin, KRKA)
		Storage:	Normal temperature
		Remarks:	Consult <i>Radio Medical</i>

Urinvejsinfektioner	7.16	Lægemiddelstof:	Pivmecillinam
		Form:	Tablet 400 mg
		Virkning:	Bakteriedræbende
		Dosering:	400 mg x 3-4 gange dagligt
		Bivirkninger:	Kvalme,diarre, træthed, nældefeber, hudkløe, hududslæt, mundsår, hovedpine
		Holdbarhed:	3 år fra produktions dato (Pivmecillinamhydrochlorid, "Paranova")
		Opbevaring:	Almindelig temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Urinary tract infection	7.16	Active Pharmaceutical Ingredients:	Pivmecillinam
		Form:	Tablets 400 mg
		Effect:	Kills bacteria
		Dosage:	400 mg x 3-4 times daily
		Side-effects:	Nausea, diarrhoea, fatigue, hives, skin itching, skin rash, mouth ulcers, headache
		Validity:	3 years from production date (Pivmecillinamhydrochlorid "Paranova")
		Storage:	Normal temperature
		Remarks:	Consult <i>Radio Medical</i>
Herpes simplex Skoldkopper Herpes zoster (helvedesild)	7.17	Lægemiddelstof:	Aciclovir
		Form:	Tablet 400 mg
		Virkning:	Hæmmer virus produktionen
		Dosering:	400-800 mg x 3 gange dagligt
		Bivirkninger:	Diarré, kvalme, opkast, feber, kraftesløshed, træthed, hovedpine, døsighed, hudkløe, hududslæt, lysfølsom hud
		Holdbarhed:	5 år fra produktions dato (Aciclovir, STADA Nordic ApS)
		Opbevaring:	Opbevares under 25 °C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Herpes simplex Variceller Herpes zoster	7.17	Active Pharmaceutical Ingredients:	Aciclovir
		Form:	Tablets 400 mg
		Effect:	Inhibits virus production
		Dosage:	400-800 mg x 3 times daily
		Side-effects:	Diarrhoea, nausea, vomiting, fever, weakness, fatigue, headache, drowsiness, itchy skin, skin rash, photosensitive skin
		Validity:	5 years from productions date (Aciclovir, STADA Nordic ApS)
		Storage:	Do not store above 25°C
		Remarks:	Consult <i>Radio Medical</i>

Grp. 8 Malaria – forebyggelse / Malaria – Prevention

Forebyggelse af malaria	8.1	Lægemiddelstof:	Atovaquon/Proguanil (Malarone®).
		Form:	Tabletter à 250/100 mg.
		Virkning:	Forebygger malaria.
		Dosering:	Tabletterne skal indtages med fedtholdigt mad (mælk/smør). Forebyggelse fra 1 dag før ankomst til 7 dage efter afrejse fra malariaområdet. Dosis voksne: 1 tablet dagligt.
		Bivirkninger:	Hovedpine, mavekneb, kvalme, opkastning, appetitløshed, diarré, hoste og allergisk hududslæt. Ved opkastning inden én time efter indtagelse af medikamentet indtages ny tablet.
		Holdbarhed:	5 år fra produktionsdato (Malarone®, GlaxoSmithKline Pharma A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Forebyggelse: Se Søfartsstyrelsens gældende vejledning om Malaria.
Prevention of malaria	8.1	Active Pharmaceutical Ingredients:	Atovaquone/Proguanil (Malarone®).
		Form:	Tablets of 250/100 mg.
		Effect:	Prevents malaria.
		Dosage:	The tablets <i>must</i> be taken with fatty food (milk/butter). The prevention starts 1 day before arrival and lasts until 7 days after departure from the malaria area. Dose, adults: 1 tablet a day.
		Side-effects:	Headache, stomach pains, nausea, vomiting, loss of appetite, diarrhoea, coughing and allergic rash. If the patient vomits within 1 hour after have taken the medicament, a new tablet has to be consumed.
		Validity:	5 years from the date of production (Malarone®, GlaxoSmithKline Pharma A/S).
		Storage:	Normal temperature.
		Remarks:	Primary prevention of malaria: See the Danish Maritime Authority's guide on Malaria.
Forebyggelse af malaria Anvendes til forebyggelse, dersom man ikke tåler 8.1	8.2	Lægemiddelstof:	Doxycyklin.
		Form:	Tabletter à 100 mg
		Virkning:	Forebygger malaria.
		Dosering:	Dosis voksne og børn over 12 år: ½ tablet dagligt. Indtages med et glas vand eller et let måltid uden mælkeprodukter. Behandlingen udstrækkes fra 1 dag før ankomst til 4 uger efter afrejse fra malariaområdet.
		Bivirkninger:	Kvalme, vaginal svamp og toksiske hudreaktioner i forbindelse med sollys.
		Holdbarhed:	3 år fra produktionsdato (Vibradox®, Sandoz).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes af gravide eller ammende eller af børn under 12 år. Direkte sollys bør undgås. Interaktion med mange lægemidler.

Grp. 8 Malaria – forebyggelse / Malaria – Prevention
(continued)

Prevention of malaria To be used for prevention if 8.1 is not tolerated	8.2	Active Pharmaceutical Ingredients:	Doxycycline
		Form:	Tablets of 100 mg.
		Effect:	Prevents malaria.
		Dosage:	To be taken with a glass of water or a light meal without any milk products. The prevention starts 1 day before arrival and lasts until 4 weeks after departure from the malaria area. Dose, adult and children over the age of 12 or 45 kg: ½ tablet daily.
		Side-effects:	Nausea, vaginal candidiasis, toxic skin reaction in connection with sunlight.
		Validity:	5 years from the date of production (Vibradox®, Sandoz).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used by pregnant or breast-feeding woman or children below the age of 12. Direct sunlight should be avoided. Interaction with many other medicines.

Grp. 8 Malaria – behandling / Malaria – Treatment
(continued)

Til behandling af P. Falciparum malaria. På baggrund af: Positiv malaria test for P. Falciparum eller almen sygdomsfornemmelse og feber kombineret med relevant risiko for malaria	8.4	Lægemiddelstof:	Artemether og lumefantrine
		Form:	Tabletter á 20mg/ 120mg
		Virkning:	Dræber malaria parasitterne
		Dosering:	Efter kontakt med <i>Radio Medical</i> Start omgående med 4 tabletter Derefter: 4 tabletter 8 timer efter første dosis 4 tabletter 24 timer efter første dosis 4 tabletter 36 timer efter første dosis 4 tabletter 48 timer efter første dosis 4 tabletter 60 timer efter første dosis
		Bivirkninger:	Mave-Tarm gener, hudkløe, hududslæt, hovedpine og svimmelhed
		Holdbarhed:	2 år fra produktionsdato (Riamet®, Novatis)
		Opbevaring:	Må ikke opbevares over 30°C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> uanset resultatet af malariatest
Treatment of P. Falciparum malaria Based on: Positive Malaria Test for P. Falciparum or general malaise and fever combined with relevant risk of malaria	8.4	Active Pharmaceutical ingredients:	Artemether and lumefantrine
		Form:	Tablets of 20mg/ 120mg
		Effect:	Kills the malaria parasite
		Dosage:	Consult <i>Radio Medical</i> 4 tablets as soon as possible Then: 4 tablets 8 hours after the first dose 4 tablets 24 hours after the first dose 4 tablets 36 hours after the first dose 4 tablets 48 hours after the first dose 4 tablets 60 hours after the first dose
		Side-effects:	Gastrointestinal disorders, itchy skin, skin rash, headache and dizziness
		Validity:	2 years from production date (Riamet®, Novatis)
		Storage:	Do not store above 30°C
		Remarks:	Consult <i>Radio Medical</i> regardless of the result of the malaria test

Grp. 9 Hjerteog kredsløb / Cardiac and circuitry

Hjertekrampe Smerter i hjertet Blodprop i hjertet Førstehjælp: Hjertestilling (elevator overkrop) 2 - 3 l ilt/minut Glycerylnitrat (9.1) Acetylsalicylsyre (9.3) Evt. for smerter: Morfin (3.4) / Tramadol (3.5)	9.1	Lægemiddelstof:	Glycerylnitrat
		Form:	Mundhule spray med 0,4 mg/dosis
		Virkning:	Udvider blodkar. Virker inden 1 minut Virkningsvarighed: 10 - 30 minutter
		Dosering:	Medikamentet pustes ind under tungen eller på kindens inderside 1 pust så hurtigt som muligt efter smerter opståen Kan gentages 1 gang Hvis smerterne ikke fortager sig, suppleres med morfin (3.4)
		Bivirkninger:	Ansigtssrødme, hovedpine og kvalme. Kan sænke blodtrykket lidt.
		Holdbarhed:	2 år fra produktionsdato (Glytrin®, Meda).
		Opbevaring:	Almindelig temperatur. Beskyttes mod frost, direkte varme og sollys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Overvåg blodtrykket, som helst ikke må komme under 90 mmHg systolisk. Må ikke anvendes i nærheden af åben ild eller gløder.
Angina pectoris Heart pain Coronary thrombosis First aid: Sit up in comfortable position 2-3 l oxygen/minute Glyceryl trinitrate (9.1) Acetylsalicylic acid (9.3) Possibly against pain: Morfin (3.4) / Tramadol (3.5)	9.1	Active Pharmaceutical Ingredients:	Glyceryl trinitrate
		Form:	Oral cavity spray with 0.4 mg/dose.
		Effect:	Expands the blood vessels. Takes effect within 1 minute. Effective for 10-30 minutes.
		Dosage:	The medicament is sprayed under the tongue or at the inner side of the cheek. 1 spray as soon as possible after the pain has occurred. May be repeated once. If the pain does not decrease, supplement with morphine (3.4).
		Side-effects:	Blushing, headache, and nausea. May cause a minor fall in blood pressure.
		Validity:	2 years from the date of production (Glytrin®, Meda).
		Storage:	Not to be stored above 25°C. Protect from frost, direct heat and sunlight.
		Remarks:	Consult <i>Radio Medical</i> . Control the blood pressure, which should not be below 90 mmHg systolic. Not to be used close to open fire or embers.

Hjertekrampe Smerter i hjertet Blodprop i hjertet Blodfortynding Se også 9.1	9.3	Lægemiddelstof:	Acetylsalicylsyre
		Form:	Tabletter à 150 mg.
		Virkning:	Virker blodfortyndende. Forhindrer blodpropper.
		Dosering:	Akut: 300 mg som tygges. Vedligehold: 75 mg dagligt.
		Bivirkninger:	Øget blødningsrisiko. Risiko for allergiske reaktioner. Farlig dosis: 150 mg per kg legemsvægt.
		Holdbarhed:	5 år fra produktionsdato (Hjertemagnyl®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke gives til patienter med mavesår.
Cardiospasm Heart pain Coronary thrombosis Blood-thinning See also 9.1	9.3	Active Pharmaceutical Ingredients:	Acetylsalicylic acid
		Form:	Tablets of 150 mg.
		Effect:	Dilute the blood. Prevents thrombosis.
		Dosage:	Acute: 300 mg has to be chewed. Hereafter: 75 mg, daily.
		Side-effects:	Increased risk of bleeding. Risk of allergic reactions. Dangerous dose: 150 mg per kg
		Validity:	5 years from the date of production (Hjertemagnyl®, Nycomed Denmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be given to ulcer patients.
Vand i lungerne Forhøjet blodtryk	9.4	Lægemiddelstof:	Furosemid.
		Form:	Tabletter à 40 mg.
		Virkning:	Tvinger væske ud af blodbanen via nyrerne til urinblæren, hvorefter væske suges fra lungerne (og andet væv). Virker 1/2 time efter indtagelse. Maksimumeffekt efter 2 timer. Virkningsvarighed: 3 - 6 timer.
		Dosering:	Vand i lungerne: 40 - 120 mg. Kan gentages efter 30 minutter. Blodtryksforhøjelse: 40 - 80 mg x 1 - 3 dagligt. (maksimalt 1500 mg/døgn).
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	5 år fra produktionsdato (Furix®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .

Lung edema High blood pressure	9.4	Active Pharmaceutical Ingredients:	Furosemide
		Form:	Tablets of 40 mg.
		Effect:	Extracts fluids from the blood vessel via the kidneys to the urinary bladder, after which the fluid is absorbed from the lungs (and other tissue). Takes effect after 1/2 hour. Maximum effect after 2 hours. Effective for 3-6 hours.
		Dosage:	Lungedema: 40-120 mg. May be repeated after 30 minutes High blood pressure: 40-80 mg once to three times daily (maximum 1500 mg/day).
		Side-effects:	None relevant.
		Validity:	5 years from the date of production (Furix®, Nycomed Denmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Vand i lungerne Forhøjet blodtryk MFAG tabel 2, 9	9.5	Lægemiddelstof:	Furosemid
		Form:	Injektionsvæske 10 mg/ml. Ampuller à 4 ml til intramuskulær eller intravenøs injektion.
		Virkning:	Tvinger væske ud af blodbanen via nyrerne til urinblæren, hvorefter væske suges fra lungerne (og andet væv). Virker inden for 5 minutter efter IV. injektion. Maksimumeffekt inden 30 minutter.
		Dosering:	Individuelt. 40 - 120 mg. Kan gentages efter 30 minutter.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	5 år fra produktionsdato (Furix®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Lung edema High blood pressure MFAG table 2, 9	9.5	Active Pharmaceutical Ingredients:	Furosemide
		Form:	Injection liquid 10 mg/ml. Ampoules of 4 ml for intramuscular or intravenous injection.
		Effect:	Extracts fluids from the blood vessel via the kidneys to the urinary bladder, after which the fluid is absorbed from the lungs (and other tissue). Takes effect 5 minutes after I.V injection. Maximum effect within 30 minutes.
		Dosage:	Dependent on the patient. 40-120 mg. May be repeated after 30 minutes.
		Side-effects:	No relevant ones.
		Validity:	5 years from the date of production (Furix®, Nycomed Denmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Forhøjet blodtryk Hjertekrampe Smerter i hjertet Forebyggelse efter blodprop i hjertet	9.6	Lægemiddelstof:	Atenolol
		Form:	Tabletter à 50 mg.
		Virkning:	Sænker puls og blodtryk.
		Dosering:	Forhøjet blodtryk: Voksne. 50-100 mg i døgnet. Hjertekrampe og hjerterytmeforstyrrelser: Sædvanligvis 50 mg 1 gang i døgnet stigende til 100 mg i døgnet. Forebyggelse efter blodprop i hjertet: 50-100 mg 1 gang i døgnet under kontrol af puls og blodtryk.
		Bivirkninger:	Træthed. Kvalme, opkastning, diarré, forstoppelse. Langsom puls, sammentrækning af blodkarrene, lavt blodtryk. Muskelsvaghed. Øget svedtendens.
		Holdbarhed:	5 år fra produktionsdato (Atenolol, Orifarm Generics A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes til patienter med astma, kroniske obstruktiv lungesygdom (KOL) og ved svær regulerbar sukkersyge.
High blood pressure Angina pectoris Heart pain Preventiv treatment after Coronary thrombosis	9.6	Active Pharmaceutical Ingredients:	Atenolol
		Form:	Tablets of 50 mg.
		Effect:	Reduces pulse and blood pressure.
		Dosage:	High blood pressure: Adults: 50- 100 mg daily Angina pectoris and heart rhythm disturbances: Usually 50 mg once a day, increasing to 100 mg daily. Prevention after coronary thrombosis: 50-100 mg once a day under the control of heart rate and blood pressure.
		Side-effects:	Fatigue, nausea, vomiting, diarrhoea, constipation, low pulse, low blood pressure, contraction of blood vessels, muscle weakness, increased sweating.
		Validity:	5 years from the date of production (Atenolol, Orifarm Generics A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Should not be used in patients with asthma, chronic obstructive pulmonary disease (COPD) and in severe controllable diabetes.

Blodfortyndende	9.8	Lægemiddelstof:	Apixaban
		Form:	Tablet á 5 mg
		Virkning:	Virker blodfortyndende. Forhindrer blodpropper
		Dosering:	5 - 10 mg x 2 dagligt, efter ordination af <i>Radio Medical</i>
		Bivirkninger:	Øget blødningstendens, kvalme hududslæt, blodig urin, vaginal blødning, blødning under huden
		Holdbarhed:	3 år fra produktion dato (Blister pk., Teva GmbH) 5 år fra produktions dato (Beholder, Teva GmbH)
		Opbevaring:	Normal temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Blood diluting	9.8	Active Pharmaceutical Ingredients:	Apixaban
		Form:	Tablets 5 mg
		Effect:	Acts as a blood thinner. Prevents blood clots
		Dosage:	5-10 mg x 2 time daily, after prescription from <i>Radio Medical</i>
		Side-effects:	Increased tendency to bleed, nausea skin rash, bloody urine, vaginal bleeding, bleeding under the skin
		Validity:	3 years from production date (Blister pack, Teva GmbH) 5 years from production date (Container, Teva GmbH)
		Storage:	Normale temperature
		Remarks:	Consult <i>Radio Medical</i>
Stopper blødninger	9.9	Lægemiddelstof:	Tranexamsyre
		Form:	Injektionsvæske 100 mg/ml
		Virkning:	Forebyggelse og behandling af blødninger
		Dosering:	1 g i.v., 3 gange dagligt
		Bivirkninger:	Diarré, kvalme, opkast, hovedpine, allergisk hudreaktion
		Holdbarhed:	3 år fra produktions dato (Tranexamsyre, Pfizer ApS)
		Opbevaring:	Normal temperatur. Må ikke fryses
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Stops Bleedings	9.9	Active Pharmaceutical Ingredients:	Tranexamicid
		Form:	Injection liquid 100 mg/ml
		Effect:	Prevention and treatment of bleedings
		Dosage:	1g i.v., 3 times daily
		Side-effects:	Diarrhoea, nausea, vomiting, headache, allergic skin reaction
		Validity:	3 years from production date (Tranexamicid, Pfizer ApS)
		Storage:	Normale temperature. Do not freeze
		Remarks:	Consult <i>Radio Medical</i>

Grp. 10 Hudmidler / Dermatological agents

Svamp Hudsvamp (Se også 14.1)	10.2	Lægemiddelstof:	Miconazol
		Form:	Crema 20 mg/g (2%).
		Virkning:	Dræber svampe i huden.
		Dosering:	Den angrebne hud indsmøres 2 gange dagligt. Behandlingen bør fortsætte 1 uge efter symptomfrihed.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	2 år fra produktionsdato (Brentan®, McNeil Denmark ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Bør ikke anvendes i åbne sår, i hårbund, skæg eller negle.
Fungus Skin fungus (See also 14.1)	10.2	Active Pharmaceutical Ingredients:	Miconazole
		Form:	Lotion 20 mg/g (2%).
		Effect:	Restricts fungus growth in the skin.
		Dosage:	Apply to the skin twice daily. The treatment should be continued for a week after the symptoms have disappeared.
		Side-effects:	No relevant ones.
		Validity:	2 years from the date of production (Brentan®, McNeil Denmark ApS).
		Storage:	Not above 25°C.
		Remarks:	Should not be used in open wounds, on scalps, beards, or nails.
Lus Fladlus Fnat	10.5	Lægemiddelstof:	Permethrin
		Form:	Crema 50 mg/g (5%).
		Virkning:	Gift, der dræber lus, fnatmider og deres æg.
		Dosering:	Påsmøres den rene og tørre hud.
		Bivirkninger:	Kløe og rødme af huden.
		Holdbarhed:	5 år fra produktionsdato (Nix®, ACO Hud Nordic AB).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Se instruktionen i indlægsseddel.
Lice Crab lice Scabies	10.5	Active Pharmaceutical Ingredients:	Permethrin
		Form:	Lotion 50 mg/g (5%).
		Effect:	Poison that kills lice, scab mites, and their eggs.
		Dosage:	Apply to clean, dry skin.
		Side-effects:	Itching and skin blushing.
		Validity:	5 years from the date of production (Nix®, ACO Hud Nordic AB).
		Storage:	Normal temperature.
		Remarks:	See package leaflet.

Grp. 10 Hudmidler / Dermatological agents
(continued)

Lus Fladlus Fnat	10.6	Lægemiddelstof:	Permethrin
		Form:	Shampo 10 mg/g (1%).
		Virkning:	Gift, der dræber lus, fnatmider og deres æg.
		Dosering:	Indgnides i det renvaskede, froterede og fugtige hår og hårbund. Shampooen skal sidde i 10 minutter før den skylles
		Bivirkninger:	Kløe og rødme af huden.
		Holdbarhed:	2 år fra produktionsdato (Nix®, ACO Hud Nordic AB).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Se instruktionen i indlægsseddel.
Lice Crab lice Scabies	10.6	Active Pharmaceutical Ingredients:	Permethrin
		Form:	Shampoo 10 mg/g (1%).
		Effect:	Poison that kills lice, scab mites, and their eggs.
		Dosage:	Apply to washed, rubbed, and wet hair and scalp. The shampoo should not be rinsed off until after 10 minutes. Dead lice and eggs are removed with a fine comb while the
		Side-effects:	Itching and skin blushing.
		Validity:	2 years from the date of production (Nix®, ACO Hud Nordic AB).
		Storage:	Normal temperature.
		Remarks:	See package leaflet.
Eksem Psoriasis Hudkløe Solskoldning Hudsvie efter insektbid	10.7	Lægemiddelstof:	Glukokortikoid gruppe II
		Form:	Crema til udvortes brug.
		Virkning:	Modvirker betændelsestilstande, der er fremkaldt af allergiske og toksiske årsager. Virker kløestillende og smertestillende.
		Dosering:	Voksne: Påsmøres tyndt 1-2 gange daglig.
		Bivirkninger:	Hudkløe brændende fornemmelse i huden, tynd hud, kontakteksem.
		Holdbarhed:	3 år fra produktionsdato (Locoid, Astellas Pharma A/S).
		Opbevaring:	Almindelig temperatur. Må ikke opbevares koldt eller fryses.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes i åbne sår.
Eczema Psoriasis Itching Sunburns Burning sensation after insect bites	10.7	Active Pharmaceutical Ingredients:	Glucocorticoid group II
		Form:	Lotion for external use.
		Effect:	Eases inflammations caused by allergic or toxic agents. Antipruritic and analgesic.
		Dosage:	Adults: Apply a thin layer to the skin 1-2 times daily.
		Side-effects:	Pruritic, burning sensation in the skin, thin skin, contact dermatitis.
		Validity:	3 years from the date of production (Locoid, Astellas Pharma A/S).
		Storage:	Normal temperature. Do not refrigerate or freeze.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used in open wounds

Grp. 10 Hudmidler / Dermatological agents
(continued)

Hudinfektioner	10.8	Lægemiddelstof:	Fusidin
		Form:	Salve 2%
		Virkning:	Bakteriedræbende
		Dosering:	Påsmøres det berørte hudområde 2-3 gange dagligt
		Bivirkninger:	Kontakteksem
		Holdbarhed:	2 år efter produktion dato (Fusidinsyre, Orifarm) 4 uger efter anbrud
		Opbevaring:	Normal temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> Må ikke anvendes i åbne sår
Skininfections	10.8	Active Pharmaceutical Ingredients:	Fusidine
		Form:	Ointment 2%
		Effect:	Kills bacteria
		Dosage:	Applies the affected skin area 2-3 times daily
		Side-effects:	Contact eczema
		Validity:	2 years from production date (Fusisdin acid, Orifarm) 4 weeks after opening
		Storage:	Normale temperature
		Remarks:	Consult <i>Radio Medical</i> Do not use in open wounds

Grp. 11 Desinficerende midler / Disinfectants

Rengøring af: 1. Instrumenter FØR og EFTER brug 2. Hænder 3. Intakt hud	11.1	Lægemiddelstof:	Klorhexidin/ethanol.
		Form:	Væske 0,5-5% / 70-85%
		Virkning:	Desinficerende
		Dosering:	-
		Anmærkninger:	Må ikke anvendes i sår, i øjne eller på slimhinder. Indeholder 85 % alkohol. Se Søfartsstyrelsens Lægebog. Alternativt indkøbes Chlorhexidin 0,5 % og Ethanol minimum 70 %, i 2 separate flasker. Først anvendes chlorhexidin, derefter ethanol.
Disinfection of: 1. Instruments BEFORE and AFTER use 2. Hands 3. Intact skin	11.1	Active Pharmaceutical Ingredients:	Chlorhexidine/ethanol
		Form:	Liquid 0.5-5% / 70-85%
		Effect:	Disinfective
		Dosage:	-
		Remarks:	Not be used in wounds, in eyes, or on mucous membranes. Contains 85% alcohol. See the Danish Maritime Authority's Medical Manual. Alternatively, Chlorhexidine 0.5% and ethanol with a minimum of 70% may be purchased in 2 separate bottles. Use chlorhexidine first and then ethanol.
Førstehjælpsbehandling af urene sår Mundskylning	11.2	Lægemiddelstof:	Klorhexidin
		Form:	Væske 0,2%
		Virkning:	Desinficerende.
		Dosering:	Kan anvendes i forurenede sår, der er utilgængelig for normal sårrensning.
		Anmærkninger:	Se Lægebog for søfarende
First aid treatment of dirty wounds Mouth rinse	11.2	Active Pharmaceutical Ingredients:	Chlorhexidine
		Form:	Liquid 0.2%
		Effect:	Disinfective.
		Dosage:	May be used for dirty wounds that are not accessible for normal wound disinfection.
		Remarks:	See the Medical guide for seafarers

Grp. 12 Midler mod mave- og tarmsygdomme / Gastro intestinal agents

Diarré	12.6	Lægemiddelstof:	Loperamid
		Form:	Tabletter à 2 mg.
		Virkning:	Hæmmer tarmenes bevægelser.
		Dosering:	Start: 2 tabletter. Pause 1 time. Derpå 1 tablet for hver uformet afføring. Maksimum: 8 tabletter (16 mg) per døgn.
		Bivirkninger:	Mundtørhed, kvalme, mavesmerter. Risiko for tarmstop.
		Holdbarhed:	5 år fra produktionsdato (Imodium®, McNeil).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Undersøg om den syge har feber. Bør kun anvendes til børn efter <i>Radio Medicals</i> anvisninger. Bør kun anvendes kortvarigt ved akut diarré hos i øvrigt upåvirkede personer.
Diarrhoea	12.6	Active Pharmaceutical Ingredients:	Loperamide
		Form:	Tablets of 2 mg.
		Effect:	Restricts the motility of the intestines.
		Dosage:	Start: 2 tablets. Wait for 1 hour. 1 tablet for every loose stool. Maximum: 8 tablets (16 mg) daily.
		Side-effects:	Dry mouth, nausea, slight stomach pains. Risk of intestinal stop.
		Validity:	5 years from the date of production (Imodium®, McNeil).
		Storage:	Normal temperature.
		Remarks:	Examine whether the patient has a fever. Should only be given to children on directions given by <i>Radio Medical</i> . Use only for a short period of time in case of acute diarrhoea on patients not otherwise affected.
Væsketab Salttab Febertilstande Opkastninger Diarré MFAG tabel 8, 10, 11.	12.7	Lægemiddelstof:	Glucose/kaliumklorid/magnesiumsulfat/natriumcitrat/natriumklorid.
		Form:	Afdelte pulvere.
		Virkning:	Erstatter væske og salte.
		Dosering:	1 pulver opløses i 1 liter kogt, afkølet vand. Basisbehov: 2,5 liter per døgn. For hver grad over 37° C gives yderligere 300 ml per døgn. For hver diarré gives yderligere 200 ml per døgn.
		Bivirkninger:	Ingen.
		Holdbarhed:	2 år fra produktionsdato (Revolyt®, Gunnar Kjems APS).
		Opbevaring:	Almindelig temperatur. Færdigblandet opløsning opbevares i
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .

Loss of fluid Loss of salt Feverish conditions Vomiting Diarrhoea MFAG tables 8, 10, 11	12.7	Active Pharmaceutical Ingredients:	Glucose/potassium/magnesium/sodium citrate/sodium
		Form:	Divided powder.
		Effect:	Replaces liquids and salts.
		Dosage:	1 powder is dissolved in 1 litre of boiled, cooled water. Basic need: 2.5 litres daily. For every degree above 37°C, the patient is given another 300 ml daily. For every instance of
		Side-effects:	None.
		Validity:	2 years from the date of production (Revolyt®, Gunnar Kjems
		Storage:	Normal temperature. The mixed dissolution should be stored in a cold place.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
Mavesår	12.8	Lægemiddelstof:	Omeprazol
		Form:	Tabletter eller kapsler 20 à mg.
		Virkning:	Hæmmer dannelsen af mavesyre.
		Dosering:	Mavesår: 20 – 40 mg dagligt. Langtidsbehandling: 10 – 20 mg dagligt.
		Bivirkninger:	Kvalme, opkastning, diarré, forstoppelse, luft i tarmen. Hovedpine, træthed, svimmelhed, søvnforstyrrelser.
		Holdbarhed:	4 år fra produktionsdato (Omeprazol, Copyfarm A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Kan eventuelt kombineres med antibiotika.
Stomach ulcer	12.8	Active Pharmaceutical Ingredients:	Omeprazol
		Form:	Tablet or capsules of 20 mg.
		Effect:	Inhibits gastric acid secretion.
		Dosage:	Stomach ulcer: 20-40 mg daily. Long term treatment: 10-20 mg daily.
		Side-effects:	Nausea, vomiting, diarrhoea, constipation, flatulence, headache, fatigue, dizziness, sleep disturbances.
		Validity:	4 years from the date of production (Omeprazol, Copyfarm A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Can be combined with antibiotics.

Sure opstød Halsbrand	12.9	Lægemiddelstof:	Antacida
		Form:	Tabletter eller oral opløsning.
		Virkning:	Neutraliserer mavesyre.
		Dosering:	Jf. indlægsseddel i præparatet. Efter ordination af <i>Radio Medical</i> .
		Bivirkninger:	Forstoppelse, diarré.
		Holdbarhed:	5 år fra produktionsdato (Alminox®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Acid indigestion Heartburn	12.9	Active Pharmaceutical Ingredients:	Antacid
		Form:	Tablet or oral solution.
		Effect:	Neutralizes acid in the stomach.
		Dosage:	See patients information leaflet. Prescription from <i>Radio Medical</i> .
		Side-effects:	Constipation, diarrhoea.
		Validity:	5 years from the date of production (Alminox®, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Hæmorider Kløe ved endetarmen Smerter ved afføring	12.10	Lægemiddelstof:	Antihæmoride creme (skal indeholde glucokortikoid og smertestillende middel)
		Form:	Salve eller creme.
		Virkning:	Lindring af smerter, hævelse, brændende fornemmelse og kløe i forbindelse med hæmoroider og analrifter.
		Dosering:	Voksne: Smøres på 2-3 gange daglig, samt efter afføring.
		Bivirkninger:	Anal slimhindeatrofi.
		Holdbarhed:	3 år fra produktionsdato (Proctosedyl, 2care4 ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Haemorrhoids Itching at the rectum Pain during defecation	12.10	Active Pharmaceutical Ingredients:	Anti haemorrhoid ointment (Must contain glucocorticoid and analgesic)
		Form:	Ointment.
		Effect:	Relief of pain, swelling and burning sensation / itching associated with haemorrhoids, anal lacerations.
		Dosage:	Apply 2-3 times daily and after defecation.
		Side-effects:	Thin skin in the anus.
		Validity:	3 years from the date of production (Proctosedyl, 2care4 ApS).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Kvalme Opkastninge MFAG tabel 7, 8, 10, 13, 15, 20	12.13	Lægemiddelstof:	Ondansetron
		Form:	Tablet á 4 mg
		Virkning:	Modvirker kvalme og opkast
		Dosering:	Efter ordination fra <i>Radio Medical</i>
		Bivirkninger:	Hovedpine, forstoppelse, rødme og varmekølelse
		Holdbarhed:	5 år fra produktions dato (Aurobindo Pharma)
		Opbevaring:	Normal temperatur
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Nausea Vomiting MFAG tabel 7, 8, 10, 13, 15, 20	12.13	Active Pharmaceutical Ingredients:	Ondansetron
		Form:	Tablets of 4 mg
		Effect:	Prevent nausea and vomiting
		Dosage:	By prescription from <i>Radio Medical</i>
		Side-effects:	Headache, Constipation, redness and warm feeling
		Validity:	5 years from production date (Aurobindo Pharma)
		Storage:	Normale temperature
		Remarks:	Consult <i>Radio Medical</i>
Forstoppelse	12.14	Lægemiddelstof:	Kaliumchlorid, Macrogol 3350, Natriumchlorid, Natriumhydrogencarbonat.
		Form:	Pulver til oral opløsning
		Virkning:	Binder væske i tarmen, hvorved afføringen blødgøres
		Dosering:	Efter ordination fra <i>Radio Medical</i>
		Bivirkninger:	Mavesmerter, Anal irritation, Diarré, Flatulens, Kvalme, Meteorisme, Opkastning
		Holdbarhed:	3 år fra produktions dato (Movicol®, Norgine BV)
		Opbevaring:	Ikke over 25°C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Constipation	12.14	Active Pharmaceutical Ingredients:	Potassium chloride, Macrogol 3350, Sodium chloride, Sodium bicarbonate
		Form:	Powder for oral solution
		Effect:	Binds fluid in the intestine, thereby softening the stools
		Dosage:	By prescription from <i>Radio Medical</i>
		Side-effects:	Abdominal pain, Anal irritation, Diarrhea, Flatulence, Nausea, Meteorism, Vomiting
		Validity:	3 years from production date (Movicol® ,Norgine BV)
		Storage:	Not above 25°C
		Remarks:	Consult <i>Radio Medical</i>

Grp. 13 Øjen- og øremidler / Agents for Eyes and Ears

Allergiske øjensygdomme Inflammatoriske øjensygdomme Øregangseksem	13.3	Lægemiddelstof:	Dexamethason
		Form:	Dråber à 1 mg/ml.
		Virkning:	Standser den allergiske eller inflammatoriske proces.
		Dosering:	Øjet: 1 - 2 dråber bag nederste øjenlåg hver time indtil bedring. Herefter 1 dråbe 4 - 6 gange per døgn.
		Bivirkninger:	Allergi. Risiko for maskering af infektioner, skader eller sygdomme i øjet.
		Holdbarhed:	3 år fra produktionsdato (Maxidex®, S.A. Alcon-couvreux
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.
Allergic eye diseases Inflammatory eye diseases Ear eczema	13.3	Active Pharmaceutical Ingredients:	Dexamethasone
		Form:	Eye drops of 1 mg/ml.
		Effect:	Stops the allergic or inflammatory process.
		Dosage:	The eye: 1-2 drops behind the lower eyelid every hour until improvement. Subsequently, 1 drop four-six times
		Side-effects:	Allergy. Risk of masking of infections, injuries, or diseases in the eye.
		Validity:	3 years from the date of production produktionsdato (Maxidex®, S.A. Alcon-couvreux N.V.).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.
Grøn stær: Smerter i øjet, Regnbuesyn Tågesyn	13.4	Lægemiddelstof:	Pilocarpin
		Form:	Dråber à 20 mg/ml.
		Virkning:	Nedsætter trykket i det indre øje.
		Dosering:	Hyppig drypning bag nederste øjenlåg efter ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Sammentrækning i pupillens lukkemuskel ("knappenålpupil") og hovedpine.
		Holdbarhed:	1½ år fra produktionsdato (Pilocarpin "ophta"®, Ophtha A/S) Må højst anvendes 1 måned efter åbning.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.

Glaucoma: Pains in the eye, Rainbow vision, Blurred vision	13.4	Active Pharmaceutical Ingredients:	Pilocarpine
		Form:	Eye drops of 20 mg/ml.
		Effect:	Reduces pressure inside the eye.
		Dosage:	Frequent drops behind the lower eyelid after prescription from <i>Radio Medical</i> .
		Side-effects:	Spasms of the sphincter of the pupil ("pin-hole pupil") and headache.
		Validity:	1½ years from the date of production (Pilocarpin "ophtha" [®] , Ophtha A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.
Herpes Helvedesild	13.6	Lægemiddelstof:	Aciclovir
		Form:	Øjensalve à 30 mg/g.
		Virkning:	Hæmmer virus.
		Dosering:	Øjet: 1 cm salve bag nederste øjenlåg 5 gange dagligt indtil 3 døgn efter symptomfrihed.
		Bivirkninger:	Øjenirritation.
		Holdbarhed:	5 år fra produktionsdato (Zovir [®] , GlaxoSmithKline Pharma A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.
Herpes Shingles	13.6	Active Pharmaceutical Ingredients:	Acyclovir
		Form:	Eye ointment of 30 mg/g.
		Effect:	Restricts virus.
		Dosage:	The eye: Apply ointment in a layer of 1 cm behind the lower eyelid five times a day until 3 days after the patient is free of
		Side-effects:	Irritation of the eye.
		Validity:	5 years from the date of production (Zovir [®] , GlaxoSmithKline Pharma A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.

Blødgøring og løsning af ørevoks.	13.9	Lægemiddelstof:	Allantoin
		Form:	Væske.
		Virkning:	Blødgør og løsner ørevoks.
		Dosering:	10-20 dråber dryppes i øret. Et stykke vat anbringes i øret og personen lejres således at dråberne forbliver i øret. Efter 20-50 min. skylles øret med lunkent vand (37°C).
		Bivirkninger:	Se indlæggelsesseddel.
		Holdbarhed:	Se produktet (Removax®).
		Opbevaring:	Opbevares ved almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes hvis der er hul i trommehinden. Må ikke anvendes ved betændelse eller smerter i øret og ved allergi overfor indholdsstofferne.
Softens and loosens cerumen	13.9	Active Pharmaceutical Ingredients:	Allantoin
		Form:	Liquid.
		Effect:	Softens and dissolving earwax.
		Dosage:	10-20 drops in the ear. A piece of cotton wool is placed in the ear and the person is placed, so that the droplets remain in the ear. After 20-50 min rinse the ear with warm water (37°C).
		Side-effects:	See package leaflet.
		Validity:	See expire date on the product (Remowax®).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Do not use if there is a hole in the eardrum. Not for use by inflammation or pain in the ear and allergy to the ingredients.
Inflammatoriske øjensygdomme	13.11	Lægemiddelstof:	Tropicamid
		Form:	Øjendråber 1%.
		Virkning:	Udvider pupillen.
		Dosering:	Akut: 1-2 dråber 3 gange dagligt.
		Bivirkninger:	Feber, forstoppelse, hjertebanken, hallucinationer, excitation, konfusion, ansigtsrødme, akkomodationsbesvær, øjenirritation, fotofobi.
		Holdbarhed:	3 år fra produktionsdato, (Mydriacyl®, S.A. Alcon-couvreur N.V.).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Inflammatory eye diseases	13.11	Active Pharmaceutical Ingredients:	Tropicamid
		Form:	Eye drops 1%.
		Effect:	Enlarges the pupil.
		Dosage:	Acut: 1-2 drops 3 times a day.
		Side-effects:	Fever, constipation, palpitations, hallucinations, excitation, confusion, flushing, difficulties with accommodation, eye irritation, photophobia.
		Validity:	3 years from the date of production, (Mydriacyl®, S.A. Alcon-couvreur N.V.).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Øregangs eksem	13.12	Lægemiddelstof:	Ciprofloxacin / Hydrocortison
		Form:	Øredråber
		Virkning:	Behandling af øregangseksem
		Dosering:	3 dråber i berørte øre 2 gange daglig
		Bivirkninger:	Ørekløe, øresmerter, svimmelhed
		Holdbarhed:	Ubrudt 2 år fra produktionsdato (Ciflox, Novartis Healthcare A/S)
		Opbevaring:	Beskytte mod lys og ikke over 25°C. Må ikke opbevares i køleskab
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Ear eczema	13.12	Active Pharmaceutical ingredients:	Ciprofloxacin / Hydrocortisone
		Form:	Eardrops
		Effect:	Treatment of ear canal eczema
		Dosage:	3 drops in affected ear twice daily
		Side-effects:	Ear itching, earache, dizziness
		Validity:	Unbroken 2 years from production date (Ciflox, Novartis Healthcare A/S)
		Storage:	Protect from light and not more than 25°C. Do not refrigerate
		Remarks:	Consult <i>Radio Medical</i>
Øjenbetændelse	13.13	Lægemiddelstof:	Kloramfenikol
		Form:	Øjensalve 1%
		Virkning:	Bakteriedræbende
		Dosering:	Smørres 3- 6 gange dagligt i øjet
		Bivirkninger:	Øjenkløe, øjenirritation, allergisk reaktion
		Holdbarhed:	2 år fra produktion dato (Kloramfenikol, 2care4)
		Opbevaring:	Se indlægsseddel. Efter anbrud ikke over 25°C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Eye infection	13.13	Active Pharmaceutical Ingredients:	Chloramphenicol
		Form:	Eye ointment 1%
		Effect:	Kills bacteria
		Dosage:	Apply 3-6 times daily in the eye
		Side-effects:	Eye itching, eye irritation, allergic reaction
		Validity:	2 years from production date (Kloramfenikol, 2care4)
		Storage:	Se leaflet. After opening not above 25°C
		Remarks:	Consult <i>Radio Medical</i>

Svejseøjne Lokalbedøvelse af øjet	13.14	Lægemiddelstof:	Diclofenac
		Form:	Øjendråber 1 mg/ml
		Virkning:	Lokalbedøvende
		Dosering:	1 dråbe 3-5 gange dagligt
		Bivirkninger:	Irritation af øje, øjenkløe, sløret syn
		Holdbarhed:	2 år fra produktion dato (Dicloabak, 2care4)
		Opbevaring:	Ikke over 25°C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Arc eye / Welders eye Local anaesthetic of the eye	13.14	Active Pharmaceutical Ingredients:	Diclofenac
		Form:	Eye drops 1mg /ml
		Effect:	Local anaesthetic.
		Dosage:	1 drop, 3-5 time daily
		Side-effects:	Eye irritation, eye itching, blurred vision
		Validity:	2 years from production date (Dicloabak, 2 care4)
		Storage:	Not above 25°C
		Remarks:	Consult <i>Radio Medical</i>

Grp. 14 Gynækologiske midler / Gynaecological agents

Svamp i skeden Hudsvamp (se også 10.2) Neglesvamp	14.1	Lægemiddelstof:	Fluconazol
		Form:	Kapsler à 150 mg.
		Virkning:	Hæmmer svampevækst.
		Dosering:	Skedesvamp: 150 mg som éngangsdosis. Hudsvamp: 150 mg per uge i 4 - 6 uger. Neglesvamp: 150 mg per uge i 3 - 12 måneder (til en ny negl er dannet).
		Bivirkninger:	Allergi. I sjældne tilfælde alvorlig leverforgiftning. Fluconazol påvirker virkningen af adskillige andre medikamenter, herunder bl.a. blodfortyndende og beroligende midler samt p-piller.
		Holdbarhed:	5 år fra produktionsdato (Fluconazol "KRKA", KRKA Sverige AB).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes til gravide.
Vaginal fungus Skin fungus (see also 10.2) Nail fungus	14.1	Active Pharmaceutical Ingredients:	Fluconazole
		Form:	Capsule of 150 mg.
		Effect:	Restricts fungal growth.
		Dosage:	Vaginal fungus: 150 mg as a single dose. Skin fungus: 150 mg a week for 4-6 weeks. Nail fungus: 150 mg a week for 3-12 months (until a new nail has been formed).
		Side-effects:	Allergy In rare cases, serious liver poisoning. Fluconazole affects the effect of several other medicaments, including inter alia blood-thinning substances, sedatives, and birth control pills.
		Validity:	5 years from the date of production (Fluconazol "KRKA", KRKA Sverige AB).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used by pregnant women.
Blødning EFTER abort eller fødsel	14.2	Lægemiddelstof:	Methylergometrin
		Form:	Tabletter à 0,125 mg.
		Virkning:	Livmoderssammentrækkende.
		Dosering:	Efter abort eller fødsel: 0,125 - 0,250 mg 3 gange dagligt.
		Bivirkninger:	Overfølsomhed, hovedpine, mavesmerter, forhøjet blodtryk, hududslet.
		Holdbarhed:	5 år fra produktionsdato (Methergin®, Novartis).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes til gravide med blødninger.

Grp. 14 Gynækologiske midler / Gynaecological agents

(continued)

Bleeding AFTER giving birth or miscarriage	14.2	Active Pharmaceutical Ingredients:	Methylergometrine
		Form:	Tablets of 0,125 mg.
		Effect:	Contracts the uterus.
		Dosage:	After giving birth or miscarriage: 0.125-0.250 mg three times daily.
		Side-effects:	Allergy, headache, stomachache, high blood pressure, rash.
		Validity:	5 years from the date of production (Methergine®, Novartis).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used by pregnant women who are bleeding.

Grp. 15 Midler mod forkølelse og astma / Agents for common Cold and Asthma

Hostestillende	15.1	Lægemiddelstof:	Dextromethorphan
		Form:	Tabletter á 30 mg.
		Virkning:	Hostestillende.
		Dosering:	15-30 mg. 3-4 gange daglig.
		Bivirkninger:	Ufrivillige muskelsammentrækning, døsighed, svimmelhed, hallucinationer, mani.
		Holdbarhed:	3 år fra produktionsdato (Dexofan® Nycomed).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Bør ikke anvendes af børn under 6 år. Bør ikke anvendes ved samtidig behandling med 7.6 og 7.7 Metronidazol. Bør ikke anvendes ved samtidig behandling med Disulfiram (Antabus®).
Cough	15.1	Active Pharmaceutical Ingredients:	Dextromethorphan
		Form:	Tablets of 30 mg.
		Effect:	Reduces coughs.
		Dosage:	15 – 30 mg. 3-4 times daily.
		Side-effects:	Unvoluntary muscle contraction, drowsiness, dizziness, hallucinations and mania.
		Validity:	3 years from the date of production (Dexofan®, Nycomed).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Should not be used by children under the age of 6 years Should not be used during concomitant treatment with 7.6 or 7.7 Metronidazole. Should not be used during concomitant treatment with disulfiram (Antabuse®).

Grp. 15 Midler mod forkølelse og astma / Agents for common Cold and Asthma (continued)

Forkølelse Bihulebetændelse Kæbehulebetændelse Pandehulebetændelse Mellemørebetændelse	15.5	Lægemiddelstof:	Xylometazolin
		Form:	Næsedråber/næsespray. 1 mg/ml.
		Virkning:	Skaffer afløb fra mellemøret, bihuler, pandehuler og kæbehuler. Virker øjeblikkeligt og i 10-12 timer.
		Dosering:	2 pust/dråber i hvert næsebor 2 - 3 gange dagligt i maksimum 10 dage.
		Bivirkninger:	Ved længere tids anvendelse kan stoffet skade næsens slimhinder.
		Holdbarhed:	3 år fra produktionsdato (Zymelin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes ved symptomer på grøn stær (se 13.4).
Cold Sinusitis Maxillary sinusitis Frontal sinusitis Otitis media	15.5	Active Pharmaceutical Ingredients:	Xylometazoline
		Form:	Nose drops/nose spray. 1 mg/ml.
		Effect:	Opens drainage from middle ears and sinuses. Takes effect immediately and stays effective for 10-12 hours.
		Dosage:	2 sprays/drops in each nostril two-three times daily for a maximum of ten days.
		Side-effects:	If used for a long period, the substance may damage the mucous membranes of the nose.
		Validity:	3 years from the date of production (Zymelin®, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used if there are symptoms of glaucoma (see 13.4).
Førstevalg ved astma eller astmalignende vejtrækning som følge af udsættelse for røg, gasarter og dampe. MFAG table 9	15.7	Lægemiddelstof:	Salbutamol
		Form:	Inhalationsspray, 0.1 mg per dosis
		Virkning:	Løser kramper i den glatte muskulatur i luftvejene
		Dosering:	1 -2 pust. Max 4 gange dagligt
		Bivirkninger:	Muskelkramper, hurtig puls, rystelser, hovedpine
		Holdbarhed:	2 år fra produktionsdato (Aeromir®, Teva B.V.)
		Opbevaring:	Ikke over 25 °C - beskyt mod kulde og frost
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
First choice of medicament in case of asthma or asthma-like breathing caused by exposure to smoke, gases, and fumes. MFAG table 9	15.7	Active Pharmaceutical Ingredients:	Salbutamol
		Form:	Spray, 0.1 mg per dose
		Effect:	Loosens convulsions in the smooth musculature in the respiratory tract
		Dosage:	1 -2 puff. Maximum 4 times daily
		Side-effects:	Muscle cramps, fast pulse, tremors, headache
		Validity:	2 years from the date of production (Aeromir®, Teva B.V.)
		Storage:	Below 25°C - protect against cold and frost
		Remarks:	Consult <i>Radio Medical</i>

Grp. 15 Midler mod forkølelse og astma / Agents for common Cold and Asthma (continued)

Astma eller astmalignende vejtrækning som følge af udsættelse for røg, gasarter og dampe. IMFAG tabel 9	15.8	Lægemiddelstof:	Glucocorticoid til inhalation
		Form:	Spray 100-200 mikrogram/ dosis
		Virkning:	Fjerner inflammatorisk ødem og hævelse i luftvejene
		Dosering:	1 pust efter behov
		Bivirkninger:	Svamp i munden, hovedpine, hæshed, hoste og halsgener
		Holdbarhed:	3 år fra produktionsdato (AeroBec, 2care4)
		Opbevaring:	Ikke over 25°C
		Anmærkninger:	Søg samråd med <i>Radio Medical</i>
Asthma or asthma-like breathing caused by exposure to smoke, gases, and fumes. MFAG tabel 9	15.8	Active Pharmaceutical Ingredients:	Glycocorticoid for inhalation
		Form:	Spray 100-200 microgram/dosage
		Effect:	Removes inflammatory oedema and swelling of the respiratory tract
		Dosage:	1 puff as required
		Side-effects:	Fungus in the mouth, headache, hoarseness, cough and sore throat
		Validity:	3 years from the date of production (AeroBec, 2care4)
		Storage:	Below 25°C
		Remarks:	Consult <i>Radio Medical</i>

Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar			
1. valgs præparat ved: Blodtab Væsketab Forebyggelse af shock MFAG tabel 7	16.1	Lægemiddelstof:	Natriumklorid
		Form:	Infusionsvæske 9 mg/ml. Beholdere à 1000 ml til intravenøs anvendelse.
		Virkning:	Erstatter væske og salte.
		Dosering:	Normalt ikke over 500 ml per time (godt 150 dråber per minut). 20 dråber = 1 ml.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Natriumklorid isotonisk, Frenius Kabi).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved malaria og ved hjertetilfælde bør infusionshastigheden være langsom af hensyn til risikoen for udvikling af lungeødem (9.4/9.5).
First choice of medicament in case of: Loss of blood Loss of liquid Prevention of shock MFAG table 7	16.1	Active Pharmaceutical Ingredients:	Sodium chloride
		Form:	Infusion liquid 9 mg/ml. Containers of 1000 ml for intravenous use.
		Effect:	Replaces liquids and salts.
		Dosage:	Normally, no more than 500 ml per hour (about 150 drops a minute). 20 drops = 1 ml.
		Side-effects:	None relevant.
		Validity:	3 years from the date of production (Sodium chloride, Frenius Kabi).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . In case of malaria and heart attacks, the infusion speed should be slow because of the risk of developing lung oedema (9.4/9.5).
Ernæringserstatning Væsketab (Se også 12.7)	16.2	Lægemiddelstof:	Glucose og natriumklorid.
		Form:	Infusionsvæske med glucose 25 mg/ml og natriumklorid 4,5 mg/ml. Beholdere à 1000 ml til intravenøs anvendelse.
		Virkning:	Erstatter væske, glucose og salte.
		Dosering:	Normalt ikke over 500 ml per time (godt 150 dråber per minut). 20 dråber = 1 ml.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Natriumklorid-Glucose "Braun", B. Braun Medical A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved malaria og ved hjertetilfælde bør infusionshastigheden være langsom af hensyn til risikoen for udvikling af lungeødem (9.4/9.5).

Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar			(continued)
Food substitute Loss of liquids (See also 12.7)	16.2	Active Pharmaceutical Ingredients:	Glucose and sodium chloride
		Form:	Infusion liquid containing glucose 25 mg/ml and sodium chloride 4.5 mg/ml. Containers of 1000 ml for intravenous use.
		Effect:	Replaces liquids, glucose and salts.
		Dosage:	Normally, no more than 500 ml per hour (about 150 drops a minute). 20 drops = 1 ml.
		Side-effects:	None relevant.
		Validity:	3 years from the date of production (Natriumklorid-Glucose" Braun", B. Braun Medical A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . In case of malaria and heart attacks, the infusion speed should be slow because of the risk of developing lung oedema (9.4/9.5).
Væske til gennemsylning af intravenøs kanyale. Sårbehandling. Øjenbehandling	16.5	Lægemiddelstof:	Natriumklorid
		Form:	Væske 0,9% i ca. 20 ml beholder.
		Virkning:	Fjerner blod fra venekanyale, renser sår og øjne.
		Dosering:	Se instruks i Søfartsstyrelsens lægebog eller efter ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Fresenius Kabi).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Fluid for flushing intravenous needle. Treatment of wounds. Treatment of eyes	16.5	Active Pharmaceutical Ingredients:	Sodium chloride
		Form:	Fluid 0,9 % in containers of 20 ml.
		Effect:	Removes blood from intravenous needle, clean wounds and eyes.
		Dosage:	See the instructions in The Maritime Medical Manual or on prescription from <i>Radio Medical</i> .
		Side-effects:	No relevant ones.
		Validity:	3 years from the date of production (Fresenius Kabi).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar (continued)			
Vand til opløsning af 7.9 Ceftriaxon pulver	16.7	Lægemiddelstof:	Sterilt vand
		Form:	Væske i 10 ml beholder
		Virkning:	-
		Dosering:	-
		Bivirkninger:	-
		Holdbarhed:	3 år fra produktionsdato (Sterilt vand Mini-plasco®, B. Braun Melsungen AG)
		Opbevaring:	Normal temperatur
		<i>Anmærkninger:</i>	-
Water for dissolution of 7.9 Ceftriaxon powder	16.7	Active Pharmaceutical Ingredients:	Sterile water
		Form:	Liquid in 10 ml container
		Effect:	-
		Dosage:	-
		Side-effects:	-
		Validity:	3 years from the date of production (sterile water mini-plasco®, B. Braun Melsungen AG)
		Storage:	Normale temperature
		<i>Remarks:</i>	-

Fortegnelse og kontroldokument over lægemiddelstoffer

Inventory and Control Document of Active Pharmaceutical Ingredients

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 1 Ilt / Oxygen						
1.1	V03AN01	Medicinsk ilt	Gas	-	20 l / 200 bar	20 l / 200 bar
1.1	V03AN01	Oxygen, Medical	Gas		20 l / 200 bar	20 l / 200 bar
1.3	V03AN01	Medicinsk ilt passende til A.1 (se note)	Gas	-	2 x 2 l / 200 bar	0
1.3	V03AN01	Oxygen Medical suitable to A.1 (see note)	Gas	-	2 x 2 l / 200 bar	0
1.3 Meddelelser A kapitel A IX B, bilag 1 / Notice A Chapter IX B, annex 1						
Grp. 2 Lokalbedøvelse / Local Anaesthetic						
2.1(π)	N01BB02	Lidocain	Inj. Væske	10 mg / ml	5 x 20 ml	0
2.1(π)	N01BB02	Lidocaine	Inj. Liquid	10 mg / ml	5 x 20 ml	0
2.2	D04AB01	Lidocain	Gel	2%	2 x 30 g	0
2.2	D04AB01	Lidocaine	Gel	2%	2 x 30 g	0
Grp. 3 Smertestillende midler / Analgesics						
3.1	N02BE01	Paracetamol	Tablet	500 mg	200 stk.	0
3.1	N02BE01	Paracetamol	Tablets	500 mg	200 pcs.	0
3.2(π)	M01AE01	Ibuprofen	Tablet	400 mg	100 stk.	0
3.2(π)	M01AE01	Ibuprofen	Tablets	400 mg	100 pcs.	0
3.4 (π)	N02AA01	Morfin Δ	Inj. Væske	10 mg / ml	10 x 1 ml	30 x 1 ml
3.4 (π)	N02AA01	Morphine Δ	Inj. Liquid	10 mg / ml	10 x 1 ml	30 x 1 ml
3.7	M02AA15	Diclofenac	Gel	11,6 mg / g	50 g	0
3.7	M02AA15	Diclofenac	Gel	11,6 mg / g	50 g	0
3.8 (π)	M01AB05	Diclofenac	Supp.	50 mg	50 stk.	0
3.8 (π)	M01AB05	Diclofenac	Supp.	50 mg	50 pcs.	0
3.9(π)	N02AA01	Morphinhydroclorid Δ	Tablet	10 mg	30 stk	0
3.9(π)	N02AA01	Morphine hydroclorid Δ	Tablets	10 mg	30 pcs.	0
Grp. 4 Allergi, shock og søsyge / Allergy, Shock and Seasickness						
4.1(π)	C01CA24	Adrenalin	Inj. Væske	1 mg / ml	5 x 1ml	0
4.1(π)	C01CA24	Epinephrine	Inj. Liquid	1 mg / ml	5 x 1ml	0
4.2	R06AX18	Acrivastin	Kapsel	8 mg	12 pcs.	0
4.2	R06AX18	Acrivastine	Capsula	8 mg	12 pcs.	0
4.3(π)	H02AB09	Hydrocortison	Inj. Tørstof + væske	100 mg	2 hætteglas	0
4.3(π)	H02AB09	Hydrocortisonsuccinate	Inj. Subst + solvens	100 mg	2 vials	0
4.6(π)	H02AB06	Prednisolon	Tablet	25 mg	30 stk.	0
4.6(π)	H02AB06	Prednisolon	Tablets.	25 mg	30 pcs.	0
4.7(π)	R06AE03	Cyclizin Δ	Tablet	50 mg	30 stk.	0
4.7(π)	R06AE03	Cyclizine Δ	Tablets	50 mg	30 pcs.	0

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 5 Antidoter / Antidotes						
5.1(π)	V03AB15	Naloxon	Inj. Væske	0,4 mg /ml	5 x 1 ml	0
5.1(π)	V03AB15	Naloxone	Inj. Liquid	0,4 mg /ml	5 x 1 ml	0
5.2(π)	H04AA01	Glucagon "K" (se note)	Inj. Tørstof + væske	1 mg	0	0
5.2(π)	H04AA01	Glucagon "R" (see note)	Inj. Subst + solvens	1 mg	0	0
5.3(π)	A03BA01	Atropine	Inj. Væske	1mg/ml	0	15 x 1ml
5.3(π)	A03BA01	Atropine	Inj. Liquid	1mg/ml	0	15 x 1ml
5.4(π)	D11AX03	Calcium gluconate	Gel	2%	0	5 x 25 g
5.4(π)	D11AX03	Calcium gluconate	Gel	2%	0	5 x 25 g
5.5(π)	A12AA02 / A12AA06	Calcium lactogluconate	Brusetablet	500 mg	0	40 stk.
5.5(π)	A12AA02 / A12AA06	Calcium lactogluconate	Effervescent tablet	500 mg	0	40 pcs.
5.6(π)	A07BA01	Aktivt kul	Pulver	50 g	0	2 x 50 g
5.6(π)	A07BA01	Charcoal, activated	Powder	50 g	0	2 x 50 g
5.7	-	Ethyl alkohol	Væske	99.9%	0	3 x 500 ml
5.7	-	Ethyl alcohol	Liquid	99.9%	0	3 x 500 ml
5.8(π)	B02BA01	Phytomenadion	Inj. Væske	10 mg / ml	0	10 x 1 ml
5.8(π)	B02BA01	Phytomenadione	Inj. Liquid	10 mg / ml	0	10 x 1 ml
5.2: 2 sæt skal medbringes af besætningsmedlemmer med insulinkrævende sukkersyge / 2 sets to be carried by crew members with insulin dependent diabetes.						
Grp. 6 Akut sindssyge og kramper / Mental disorder and convulsion						
6.2(π)	N05BA01	Diazepam Δ	Tablet	5 mg	25 stk.	0
6.2(π)	N05BA01	Diazepam Δ	Tablets	5 mg	25 pcs.	0
6.3(π)	N05BA01	Diazepam Δ	Klysma	4 mg / ml	10 x 2½ ml	0
6.3(π)	N05BA01	Diazepam Δ	Enema	4 mg / ml	10 x 2½ ml	0
6.4(π)	N05AD01	Haloperidol Δ	Inj. Væske	5 mg / ml	10 x 1 ml	0
6.4(π)	N05AD01	Haloperidol Δ	Inj. Liquid	5 mg / ml	10 x 1 ml	0
6.5(π)	N05CF02	Zolpidem Δ	Tablet	10 mg	20 stk.	0
6.5(π)	N05CF02	Zolpidem Δ	Tablets	10 mg	20 pcs.	0

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 7 Infektioner / Infections						
7.1(π)	J01CE02	Phenoxymethylpenicillin	Tablet	660-800 mg	100 stk.	0
7.1(π)	J01CE02	Phenoxymethylpenicillin	Tablets	660-800 mg	100 pcs.	0
7.2(π)	J01CA04	Amoxicillin	Tablet	500 mg	30 stk.	0
7.2(π)	J01CA04	Amoxicillin	Tablets	500 mg	30 pcs.	0
7.6(π)	P01AB01	Metronidazol	Tablet	500 mg	24 pcs.	0
7.6(π)	P01AB01	Metronidazole	Tablets	500 mg	24 pcs.	0
7.7(π)	P01AB01	Metronidazol	Supp.	500 mg	20 stk.	0
7.7(π)	P01AB01	Metronidazole	Supp.	500 mg	20 pcs.	0
7.8(π)	J01MA02	Ciprofloxacin	Tablet	250 mg	40 stk.	0
7.8(π)	J01MA02	Ciprofloxacin	Tablets	250 mg	40 pcs.	0
7.9(π)	J01DD04	Ceftriaxon (se note)	inj. Tørstof	1 g	20 x 1g	0
7.9(π)	J01DD04	Ceftriaxon (see note)	inj. Subst	1 g	20 x 1g	0
7.10(π)	-	Tetanus holdig vaccine "K"	Inj. Væske	-	1 stk.	0
7.10(π)	-	Vaccine containing Tetanus "R"	Inj. Liquid	-	1 pc.	0
7.11(π)	J06BB02	Tetanus immunoglobulin "K"	Inj. Væske	250 IU.	1 x 1 ml	0
7.11(π)	J06BB02	Tetanus immunoglobulin "R"	Inj. Liquid	250 IU.	1 x 1 ml	0
7.12(π)	J01CF05	Flucloxacillin	Tablet	500 mg	50 pcs	0
7.12(π)	J01CF05	Flucloxacillin	Tablet	500 mg	50 pcs	0
7.15(π)	J01FA10	Azithromycin (se note)	Tablet	500 mg	5 x 3 stk.	0
7.15(π)	J01FA10	Azithromycin (see note)	Tablets	500 mg	5 x 3 pcs.	0
7.16(π)	J01CA08	Pivmecillinam	Tablet	400 mg	50 stk.	0
7.16(π)	J01CA08	Pivmecillinam	Tablets	400 mg	50 pcs.	0
7.17(π)	J05AB01	Aciclovir	Tablet	400 mg	70 stk.	0
7.17(π)	J05AB01	Aciclovir	Tablets	400 mg	70 pcs.	0
ADVARSEL: I DK anvendes Ceftriaxon (7,9) i stedet for den i MFAG angivet Cefuroxim. WARNING: In DK is Ceftriaxone (7.9) used instead of Cefuroxime as described in the MFAG. 7.15 Azithromycin erstatter den i MFAG angivet Erythromycin. 7.15 Azithromycin replace the Erythromycin specified in the MFAG. 7.9 Skibe med en overfartstid under 24 timer kan reducere mængden af Ceftriaxon fra 20 x1g til 5 x1 g. OBS skal blandes med sterilt vand 16.7 7.9 Ships which are engaged in voyage less than 24 hours may reduce the amount of Ceftriaxon from 20 x1g to 5 x1 g. OBS shall be mixed with sterile water 16.7						
Grp. 8 Malaria / Malaria (se note /see note)					Treatment	Prevention
8.1(π)	P01BB51	Atovaquon/Proguanil	Tablet	250 / 100 mg	0	24 stk./ besætningmedl
8.1(π)	P01BB51	Atovaquon/Proguanil	Tablet	250 / 100 mg	0.	24 pcs./ crewmember
8.2(π)	J01AA02	Doxycyclin	Tablet	100 mg	0	80 stk.
8.2(π)	J01AA02	Doxycycline	Tablet	100 mg	40	80 pcs.
8.4(π)	P01BF01	Artemether og lumefantrine	tbl.	20 mg / 120 mg	24 stk	0
8.4(π)	P01BF01	Artemether and lumefantrine	tablets.	20 mg / 120 mg	24 pcs	0
Malariamidler til behandling skal forefindes i skibe, der sejler til områder med malaria - uanset risiko. Medicinsk forebyggelse kræves hovedsagelig ved sejlads i høj risikoområder. Højrisikoområder findes opdateret og beskrevet i Søfartsstyrelsens malariastrategi på følgende link http://www.soefartsstyrelsen.dk. Drugs for treatment of malaria must be carried by all ships sailing to or in areas where malaria is recorded - irrespective of the risk. Medical prevention is mainly required for ships sailing in high risk areas. The High risk areas are available and updated on the "Danish Maritime Authority's malaria strategy" on the following link http://www.dma.dk						

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 9 Hjerteog kredsløb / Cardiac and circuitry						
9.1	C01DA02	Glycerylnitrat	Mund spray	0,4 mg/dose	1x200 doser	0
9.1	C01DA02	Glyceryl trinitrate	Oral spray	0,4 mg/dose	1x200 doses	0
9.3	B01AC06	Acetylsalisylsyre	Tablet	150 mg	100 stk.	0
9.3	B01AC06	Acetylsalicylic Acid	Tablet	150 mg	100 pcs.	0
9.4(π)	C03CA01	Furosemid	Tablet	40 mg	50 stk.	0
9.4(π)	C03CA01	Furosemide	Tablet	40 mg	50 pcs.	0
9.5(π)	C03CA01	Furosemid	Inj. væske	10 mg/ml	10x4 ml	0
9.5(π)	C03CA01	Furosemide	Inj. Liquid	10 mg/ml	10x4 ml	0
9.6(π)	C07AB03	Atenolol	Tablet	50 mg	100 stk.	0
9.6(π)	C07AB03	Atenolol	Tablet	50 mg	100 pcs.	0
9.8(π)	B01AF02	Apixaban	Tablet	5 mg	28 stk.	0
9.8(π)	B01AF02	Apixaban	Tablets	5 mg	28 pcs.	0
9.9(π)	B02AA02	Tranexamsyre	Inj. væske	100 mg/ml	10 x 5 ml	0
9.9(π)	B02AA02	Tranexamsyre	Inj. liquid	100 mg/ml	10 x 5ml	0
Grp. 10 Hud / Skin						
10.2(π)	D01AC02	Miconazol	Crema	2%	4 x15 g	0
10.2(π)	D01AC02	Miconazol	Lotion/Cream	2%	4 x15 g	0
10.5	P03AC04	Permethrin	Crema	5%	30 g	0
10.5	P03AC04	Permethrin	Lotion/Cream	5%	30 g	0
10.6	P03AC04	Permethrin	Shampoo	1%	50 ml	0
10.6	P03AC04	Permethrin	Shampoo	1%	50 ml	0
10.7(π)	D07AB02	Glucocorticoid gruppe II	Crema	Gruppe II	2 x 30 g	0
10.7(π)	D07AB02	Glucocorticoid group II	Lotion/Cream	Group II	2 x 30 g	0
10.8(π)	D06AX01	Fusidin	Cream / Salve	2%	4 x15 g	0
10.8(π)	D06AX01	Fusidine	Cream / Lotion	2%	4 x 15 g	0
Grp. 11 Desinficering / Desinfectants						
11.1	-	Klorhexidin/ethanol (se note)	Væske	0.5-5% / 70-85%	175 ml	0
11.1	-	Chlorhexidine/ethanol (see note)	Liquid	0.5-5% / 70-85%	175 ml	0
11.2	-	Klorhexidin	Væske	0.2 %	4 x 250 ml	0
11.2	-	Chlorhexidine	Liquid	0.2 %	4 x 250 ml	0
11.1: Alternativt kan anvendes Klorhexidin 0,5 - 5 % og Ethanol minimum 70-85 %, i 2 separate flasker. Blandes 1:1 Alternatively, Chlorhexidine 0,5 -5 % and ethanol with a minimum of 70 - 85 % in 2 separate bottles. Mix 1:1						

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 12 Mave og tarm / Gastrointestinal						
12.6	A07DA03	Loperamid	Tablet	2 mg	60 pcs.	0
12.6	A07DA03	Loperamide	Tablet	2 mg	60 pcs.	0
12.7	-	Glukose/kalium/natriumcitrat /natriumklorid	Pulver	13,5/1,5/2,9/ 2,6 g/pulver	18 x 20,5 g	0
12.7	-	Glucose/potassium/sodiumcitrate / sodiumchlorid	Powder	13,5/1,5/2,9/ 2,6 g/powder	18 x 20,5 g	0
12.8(π)	A02BC01	Omeprazol	Tablet/Kapsel	20 mg	28 pcs.	0
12.8(π)	A02BC01	Omeprazol	Tablet/Capsula	20 mg	28 pcs.	0
12.9	A02A	Antacida	Tablet eller oral opløsning	-	40 stk. eller 250 ml	0
12.9	A02A	Antacid	Tablet or oral solution		40 pcs. or 250 ml	0
12.10	C05A	Hæmoride creme (skal indeholde glukokortikoid and smertestillende)	Creme/Salve	-	2 x 30 g	0
12.10	C05A	Anti hæmorrhoid agent (Must contain glucocorticoid and analgesi)	Lotion/Ointment	-	2 x 30 g	0
12.13	A04AA01	Ondasetron (Se note)	Tablet	4 mg	10 stk.	0
12.13	A04AA01	Ondasetron (see note)	Tablets.	4 mg	10 pcs.	0
12.14	A06AD65	Kaliumchlorid, Macrogol 3350, Natriumchlorid, Natriumhydrogencarbonat.	oral opløsning	46,6 mg kaliumchlorid, 13,125 g macrogol 3350, 350,7 mg natriumchlorid , 178,5 mg natriumhydro gencarbonat.	4 bokse af 8 stk.	0
12.14	A06AD65	Potassium chloride, Macrogol 3350, Sodium chloride, Sodium bicarbonate	oral solution	46,6 mg kaliumchlorid, 13,125 g macrogol 3350, 350,7 mg natriumchlorid , 178,5 mg natriumhydro gencarbonat.	4 boxes of 8 pcs.	0
12.13 Erstatte, den i MFAG nævnte, injektion Metroclorpramid / 12.13 Replaces the injection Metroclorpramide described in MFAG.						

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A1
Grp. 13 Øjne og øre / Eyes and ears						
13.3(π)	S01BA01	Dexamethason	Dråber	1 mg / ml	5 ml	0
13.3(π)	S01BA01	Dexamethasone	Drops	1 mg / ml	5 ml	0
13.4(π)	S01EB51	Pilocarpin	Dråber	2 %	10 ml	0
13.4(π)	S01EB51	Pilocarpine	Drops	2 %	10 ml	0
13.6(π)	S01AD03	Aciclovir	Salve	3 %	4,5 g	0
13.6(π)	S01AD03	Aciclovir	Ointment	3 %	4,5 g	0
13.9	-	Medium til at blødgøre ørevoks, inkl. Øresprøjte	-	-	10 ml	0
13.9	-	Medium to soften earwax, incl. Earsyringe	-	-	10 ml	0
13.11(π)	S01FA06	Tropicamid	Dråber	1 %	15 ml	0
13.11(π)	S01FA06	Tropicamide	Drops	1 %	15 ml	0
13.12(π)	S02CA03	Ciprofloxacin / Hydrocortison	øredråber	10+2mg/ml	3 x 10 ml	0
13.12(π)	S02CA03	Ciprofloxacin / Hydrocortisone	eardrops	10+2mg/ml	3 x 10 ml	0
13.13(π)	S01AA01	Kloramfenikol	øjensalve	1%	2 x 4,3 - 5 g	0
13.13(π)	S01AA01	Chloramphenicol	eye ointment	1%	2 x 4,3 - 5 g	0
13.14(π)	S01BC03	Diclofenac	øjendråber	1 mg/ml	5 x10 ml	0
13.14(π)	S01BC03	Diclofenac	eye drops	1 mg/ml	5 x10 ml	0
Grp. 14 Gynækologi / Gynaecology						
14.1(π)	J02AC01	Fluconazol	Kapsel	150 mg	2 stk.	0
14.1(π)	J02AC01	Fluconazole	Capsula	150 mg	2 pcs.	0
14.2(π)	G02AB01	Methyleergometrin (se note)	Tablet	0,125 mg	10 pcs.	0
14.2(π)	G02AB01	Methyleergometrine (see note)	Tablet	0,125 mg	10 pcs.	0
12.8 skal kun forefindes, HVIS der er kvinder om bord. Shall only be available IF women on board.						
Grp. 15 Astma og forkølelse / Astma and common cold						
15.1(π)	R05DA09	Dextromethorphan	Tablet	30 mg	40 stk.	0
15.1(π)	R05DA09	Dextromethorphan	Tablet	30 mg	40 pcs.	0
15.5	R01AA07	Xylometazolin	Dråber / Spray	1 mg / ml	3 x 10 ml	0
15.5	R01AA07	Xylometazoline	Drops / Spray	1 mg / ml	3 x 10 ml	0
15.7(π)	R03AC02	Salbutamol	Spray	0,1 mg/dosis	200 doser	0
15.7(π)	R03AC02	Salbutamol	Spray	0,1 mg/dosage	200 dosis	0
15.8(π)	R03BA01	Glucocorticoid (inhalation)	Spray	100-200 mikrogram pr. dosis	100 doser	4 x 100 doser
15.8(π)	R03BA01	Glucocorticoid (inhalation)	Spray	100-200 mikrogram per dosage	100 dosis	4 x 100 dosis
Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar (se note / see note)						
16.1(π)	B05BB01	Natriumklorid	Inf. Væske	0.9 %	10 x 1 liter	0
16.1(π)	B05BB01	Sodium chloride	Inf. Liquid	0.9 %	10 x 1 litre	0
16.2(π)	B05BB02	Glucose + Natriumklorid	Inf. Væske	25/4,5 mg/ml	10 x 1 liter	0
16.2(π)	B05BB02	Glucose + Sodium chlorid	Inf. Liquid	25/4,5 mg/ml	10 x 1 litre	0
16.5(π)	V07AB	Natriumklorid 20 ml	Væske	0.9 %	20 x 20 ml	0
16.5(π)	V07AB	Sodium chloride 20 ml	Liquid	0.9 %	20 x 20 ml	0
16.7	V07AB	Steril vand til blanding af 7.9	Parenteral væske	-	20 x10 ml	0
16.7	V07AB	Sterile water to mix 7.9	Parenteral Liquid	-	20 x 10 ml	0
Infusionsvæskeerne må ikke være i glasbeholder Infusion liquids may not be in glass container						

Besætningens medbragte medicin / Crew member's private medicine

[illegible]

Besætningens medbragte medicin / Crew member's private medicine

[illegible]

Fortegnelse og Kontrolldokument over medicinsk udstyr

Inventory and Control document of medical equipment

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
A. Genoplivningsudstyr / Resuscitation equipment			
A.1	Mobilt genoplivningsudstyr, klargjort (se note)	1 sæt	0
A.1	Mobile resuscitation equipment (see note)	1 set	0
A.4	Tungetholdere, 80 mm	0	2 stk.
A.4	Guidel airway, 80 mm	0	2 pcs.
A.5	Tungetholdere, 90 mm	1 stk.	1 stk.
A.5	Guidel airway, 90 mm	1 pcs.	1 pcs.
A.6	Tungetholdere, 100 mm	0	2 stk.
A.6	Guidel airway, 100 mm	0	2 pcs.
A.7	Ekstra ventilationspose med ventil og voksenmaske	0	1 stk.
A.7	Extra ventilation bag with valve and mask	0	1 pcs.
A.8	Kraftig klædesaks	1 stk.	0
A.8	Strong scissors for clothes	1 pcs.	0
A.9	Sugekatetre, lige, CH 12	5 stk.	0
A.9	Suction catheters, straight, CH 12	5 pcs.	0
A.10	Tæppe til behandling af hypotermi	1 stk.	0
A.10	Blanket for treatment of hypothermia	1 pcs.	0
A.11	Almindelige tæpper	6 stk.	0
A.11	Ordinary blankets	6 pcs.	0
A.12	Saturationsmåler (fingermodel)	1 stk.	0
A.12	Puls oximeter (finger)	1 pcs.	0
A.13	Hudson maske med reservoirpose med ca. 1,8 m slange	5 stk.	5 stk.
A.13	Hudson mask with reservoir bag with approx. 1.8 m tube	5 pcs.	5 pcs.
A.18	Hjertestarter (AED)	1 stk.	0
A.18	Automatic extern defibrillator (AED)	1 pc.	0
A.19	Iltbrille	5 stk.	0
A.19	Nasal Cannula	5 pcs.	0
A.1: Meddelelser A kapitel A IX B, bilag 1 / A.1: Notice A Chapter IX B, annex 1			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
B. Forbindinger og plastre / Dressings and adhesive tape			
B.1	Sterile gazekompresser, ca.5 x 5 cm, ca.8 lag	20 stk.	0
B.1	Sterile gaze compress, approx.5 x 5 cm, approx. 8 layers	20 pcs.	0
B.2	Sterile gazekompresser, ca. 10 x 10 cm, ca. 8 lag	20 stk.	0
B.2	Sterile gaze compress, approx.10 x10 cm, approx. 8 layers	20 pcs.	0
B.3	Steril absorberende ikke-klæbende forbindelse, ca. 20 x 30 cm	5 stk.	0
B.3	Sterile absorbent non-adhesive surgical dressing bandage, approx. 20 x 30 cm	5 pcs.	0
B.4	Sterile kompresforbindinger, ca. 8 cm	10 ruller	0
B.4	Sterile compress bandages, approx. 8 cm	10 rolls	0
B.5	Imprægnerede net, ca. 10 x 10 cm, "Jelonet"	2 x 10 stk.	0
B.5	Impregnated nets, approx. 10 x 10 cm, "Jelonet"	2 x 10 pcs.	0
B.6	Steril absorberende forbindinger, ikke klæbende, ca. 10 x 10 cm	5 stk.	0
B.6	Sterile non-adhesive surgical dressing bandages, approx. 10 x 10 cm	5 pcs.	0
B.7	Gazeviskers, usterile	100 stk.	0
B.7	Gauze swaps, non-sterile	100 pcs.	0
B.8	Tubegaze til fingerforbindinger	10 stk.	0
B.8	Tube shaped gauze for finger bandaging	10 pcs.	0
B.9	Elastisk gazebind, ca. 4 cmx4m	10 ruller	0
B.9	Elastic gauze bandage, approx. 4 cmx4m	10 rolls	0
B.10	Hæfteplaster med sårpude, forskellige størrelser, lommepakning	1 pakke	0
B.10	Adhesive tape with compress, various sizes, pocket pack	1 pack	0
B.11	Hæfteplaster med sårpude, ca. 6 cm x 1m	5 pakker	0
B.11	Adhesive tape with compress, approx. 6 cm x 1m	5 packs	0
B.12	Fikseringsplaster Fixomull, ca. 10 cm x 10 m	1 rulle	0
B.12	Adhesive tape Fixomull, approx. 10 cm x 10 m	1 roll	0
B.13	Fikseringsplaster, "Englehud", ca. 2 cm	2 ruller	0
B.13	Adhesive tape, "peau d'ange", approx. 2 cm	2 rolls	0
B.18	Årepresse til massiv blødning	1 stk	0
B.18	Tourniquet for massive bleeding	1 pc.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
C. Udstyr til hygiejne / Equipment for sanitation			
C.1	Engangshandsker af latex, usterile, X-large	100 stk.	0
C.1	Disposable gloves, latex, non-sterile, X-large	100 pcs.	0
C.2	Engangshandsker af latex, usterile, large	100 stk.	0
C.2	Disposable gloves, latex, non-sterile, large	100 pcs.	0
C.5	Vaskesvampe, éngangs	25 stk.	0
C.5	Bath sponge, disposable	25 pcs.	0
C.6	Beskyttelsesbriller	1 stk.	0
C.6	Protective goggles	1 pc.	0
C.7	Steril afdækning, ca. 50 x 50 cm	5 stk.	0
C.7	Sterile drape, approx. 50 x 50 cm	5 pcs.	0
C.8	Flydende medicinsk sæbe pH 4,5	250 ml	0
C.8	Liquid medical soap pH 4,5	250 ml	0
C.9	Mundbind 3 lags, bredde 18 cm	50 stk.	0
C.9	Mask, 3 layer, width 18 cm	50 pcs.	0
C.10	Ligpose voksen	1 stk.	0
C.10	Body bag adult	1 pc.	0
C.11	Engangshandsker af latex, usterile, medium	100 stk.	0
C.11	Disposable gloves, latex, non-sterile, medium	100 pcs.	0
C.12	Engangshandsker af latex, usterile, small	100 stk.	0
C.12	Disposable gloves, latex, non-sterile, small	100 pcs.	0
C.13	Engangsforklæde	10 stk.	0
C.13	Disposable apron	10 pcs.	0
C.14	Lagen, sterilt	1 stk	0
C.14	Bed sheet, sterile	1 pc.	0
C.15	Absorberende underlag, ca.6 lag, 60x40 cm, usteril, engangs	10 stk.	0
C.15	Absorbent underlay, approx.6 layers, 60x40 cm, non-sterile, disposable	10 pcs.	0
C.16	Opkastpose	10 stk.	0
C.16	Vomit bag	10 pcs.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
D. Sårbehandling / Treatment of wounds			
D.1	Suturtape, "Steri-Strip™"	25 stk.	0
D.1	Suturetape, "Steri-Strip™"	25 pcs.	0
D.2	Klæbesutur, Dermabond	6 ampuller	0
D.2	Adhesive suture, Dermabond	6 ampoules	0
D.4	Ikke-absorberbar sutur, 3.0; 45 cm. FS eller DS 25	12 stk.	0
D.4	Non-absorbable suture, 3.0; 45 cm. FS or DS 25	12 pcs.	0
D.5	Rustfri stålæske med låg, ca. 18x9x3 cm	1 stk.	0
D.5	Instrument box, stainless steel with lid, approx. 18x9x3 cm	1 pc.	0
D.6	Engangsskalpel, steril, nr. 23	5 stk.	0
D.6	Disposable scalpel, sterile, no. 23	5 pcs.	0
D.7	Kirurgisk saks med lige spids	1 stk.	0
D.7	Surgical scissors w/straight points	1 pc.	0
D.8	Kirurgisk saks med krum spids	1 stk.	0
D.8	Surgical scissors w/curved points	1 pc.	0
D.9	Kirurgisk pincet, ca. 14 - 16 cm	1 stk.	0
D.9	Surgical forceps (tweezers), approx. 14 -16 cm	1 pc.	0
D.10	Anatomisk pincet	1 stk.	0
D.10	Anatomical forceps (tweezers)	1 pc.	0
D.11	Splintpincet	1 stk.	0
D.11	Forceps, splinter (tweezers)	1 pc.	0
D.12	Pean, krum	2 stk.	0
D.12	Pean, curved	2 pcs.	0
D.13	Nåleholder	2 stk.	0
D.13	Suture - holder	2 pcs.	0
D.14	Barbermaskine (skraber), engangs	3 stk.	0
D.14	Safety razor, disposable single use	3 pcs.	0
D.15	Absorberbar hæmostatisk gelatinesvamp, ca.1x1x1 cm	8 stk.	0
D.15	Haemostatic absorbable gelatine sponge, approx.1x1x1 cm	8 pcs.	0
D.17	Ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25	12 stk.	0
D.17	Non-absorbable suture, 5.0; 45 cm. FS eller DS25	12 pcs.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
E. Udstyr til injektion og infusion / Equipment for injection and infusion			
E.1	Kanylebøtte	1 stk.	0
E.1	Box for contaminated needles	1 pc.	0
E.2	Afspritningssvaps m. isopropylalkohol 70%	200 stk.	0
E.2	Alcohol swabs, isopropyl alcohol 70%	200 pcs.	0
E.3	Staserem	2 stk.	0
E.3	Tourniquet for i.v.	2 pcs.	0
E.4	Intravenøs kanyler, 1,1 x 32, 20 g (se note)	20 stk.	0
E.4	Intravenous cannula, 1.1 x 32, 20 g (see note)	20 pcs.	0
E.5	Fikseringsplaster til intravenøs kanyler	20 stk.	0
E.5	Fixing plaster for intravenous cannula	20 pcs.	0
E.6	Kanyler, steril til engangsbrug, 0,6 x 30 (~ 23G x 1 1/4) (se note)	50 stk.	50 stk.
E.6	Needle, sterile, disposable single use, 0.6 x 30 (~ 23G x 1 1/4) (see note)	50 pcs.	50 pcs.
E.7	Sprøjter, sterile til engangsbrug, Luer, 2 ml	20 stk.	80 stk.
E.7	Syringes, sterile, single use, Luer, 2 ml	20 pcs.	80 pcs.
E.8	Sprøjter, sterile til engangsbrug, Luer, 5 ml	20 stk.	0
E.8	Syringes, sterile, single use, Luer, 5 ml	20 pcs.	0
E.9	Sprøjter, sterile til engangsbrug, Luer, 10 ml	20 stk.	0
E.9	Syringes, sterile, single use, Luer, 10 ml	20 pcs.	0
E.10	Infusionssæt med gevindtilslutning til E.4, sterile til engangsbrug	10 stk.	0
E.10	Infusion set with threaded connection for E.4, sterile, single use	10 pcs.	0
E.4+E.6: Kanyler i kaliber 0,8 og intravenøs kanyler kaliber 1,2, som beskrevet i MFAG, er erstattet af kanyler i kaliber 0,6 og intravenøs kanyler kaliber 1,1. E.4+E.6: Needles size 0.8 and intravenous cannula size 1.2, as described in MFAG, are replaced with needles size 0.6 and intravenous cannula size 1.1			
F. Udstyr til medicinsk undersøgelse / Equipment for medical examination			
F.1	Blodtryksmåler	1 stk.	0
F.1	Instrument for measuring blood pressure	1 pc.	0
F.2	Stetoskop	1 stk.	0
F.2	Stethoscope	1 pc.	0
F.3	Digitalt lægetermometer	2 stk.	0
F.3	Digital clinical thermometer	2 pcs.	0
F.4	Pencillygte m. batterier	2 stk.	0
F.4	Pencil lamp w/batteries	2 pcs.	0
F.5	Tungespatel, usterile, engangs	10 stk.	0
F.5	Tongue depressor, non-sterile, single use disposable	10 pcs.	0
F.7	Hypotermi Termometer (se note)	1 stk.	0
F.7	Hypothermia thermometer (see note)	1 pc.	0
*) F.7 Termometret skal kunne måle minimum ned til 26°C. Bruges i endetarm. *) F.7 the thermometer have to be able to minimum measure down to 26°C. The thermometer is used rectal.			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
G. Udstyr til undersøgelse af blod og urin / Equipment for examination of blood and urine			
G.1	"Multistix® 5", urintest for blod/glukose/protein/nitrit/leukocytter	50 stk.	0
G.1	"Multistix® 5", Urinetest for blood/glucose/protein/nitrite/leukocytes	50 pcs.	0
G.2	Graviditetstest	2 stk.	0
G.2	Pregnancy test kit	2 pcs.	0
G.5	Malaria test, minimum for Plasmodium Falciparum, inkl. reagens og brugervejledning.	6 stk.	0
G.5	Malaria test, minimum for Plasmodium Falciparum, incl. reagens and userinstruction.	6 pcs.	0
G.6	Lancetter (fingerprikker)	6 stk.	0
G.6	Lancets (finger pricker)	6 pcs.	0
G.7	Udstyr til måling af blodsukker (se note)	0	0
G.7	Equipment for measuring bloodsugar (see note)	0	0
G.8	Hurtig test C-reaktiv protein (CRP)	5 stk.	0
G.8	Rapid test for C- reaktive proteiner (CRP)	5 pcs.	0
G.2 Skal kun forefindes, HVIS der er kvinder om bord. G.2 Shall only be available IF women on board. G.5+G.6 Malarietest og Lancetter skal forefindes i skibe, der sejler til områder med malaria - uanset risiko. G.5+G.6 Malaria test and Lancets must be carried by all ships sailing to or in areas where malaria is recorded - irrespective of the risk. G.7 1 sæt skal medbringes af besætningsmedlemmer med insulinkrævende sukkersyge G.7 1 set to be carried by crew members with insulin dependent diabetes.			
H. Øjenundersøgelse og behandling / Eye examination and treatment			
H.1	Pencillygte med UV lys, inkl. batterier	1 stk.	0
H.1	Lamp with UV light w/batteries	1 pc.	0
H.2	Fluoretsstrips til farvning af øjet	10 stk.	0
H.2	Fluorescent colour for the eye	10 pcs.	0
H.3	Vatpinde, sterile	50 stk.	0
H.3	Cotton sticks, sterile	50 pcs.	0
H.4	Horners hulmejsel	1 stk.	0
H.4	Horners spud	1 pc.	0
H.5	Øjenklap	2 stk.	0
H.5	Eye patch	2 pcs.	0
H.6	Øjenskyllevæske, Natriumchlorid 0,9%, 250 ml	8 stk.	0
H.6	Eye irrigating liquid, Sodium chloride 0.9%, 250 ml	8 pcs.	0
I. Udstyr til tømning af urinblæren / Equipment for drainage of urinary bladder			
I.1	Engangskatetre m. krum spids, CH 14, til mænd	10 stk.	0
I.1	Catheters for single use w/ curved tip, CH 14 for men	10 pcs.	0
I.2	Urinkelbe, plastic	1 stk.	0
I.2	Bed bottle, plastic	1 pc.	0
I.3	Engangskatetre CH 14 til kvinder	5 stk.	0
I.3	Catheters for single use, CH 14 for women	5 pcs.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
J. Udstyr til forstuvninger og brud / Equipment for sprains and fractures			
J.1	ColdHot™ kulde/varme omslag, 11 x 27 cm	1 stk.	0
J.1	ColdHot™ pack 11x24 cm	1 pc.	0
J.2	Elastikbind, ca. 10 cmx5m	3 stk.	0
J.2	Elastic bandage, approx. 10 cmx5m	3 pcs.	0
J.3	Elastikbind, ca.6 cm x 5m	1 stk.	0
J.3	Elastic bandage, approx. 6 cmx5m	1 pc.	0
J.4	Trekantet tørklæde (Mitella)	2 stk.	0
J.4	Triangular scarf (Mitella)	2 pcs.	0
J.5	Sikkerhedsnåle	12 stk.	0
J.5	Safety pins	12 pcs.	0
J.6	Fingerskinne til immobilisering af ét led	5 stk.	0
J.6	Finger splint for immobilisation of one joint	5 pcs.	0
J.7	Fingerskinne til immobilisering af to led	5 stk.	0
J.7	Finger splint for immobilisation of two joints	5 pcs.	0
J.8	Polsterbind, ca. 10 cm x 2,7 m	10 stk.	0
J.8	Padded bandages, approx. 10 cm x 2.7 m	10 pcs.	0
J.9	Gipsbind, ca. 12 cmx2m	10 stk.	0
J.9	Plaster bandage, approx. 12 cmx2m	10 pcs.	0
J.10	Fikseringsplaster, Tensoplast, ca. 2 cmx2m	2 ruller	0
J.10	Adhesive tape, Tensoplast, approx. 2 cmx2m	2 rolls	0
K. Udstyr til immobilisering og transport / Equipment for immobilisation and transportation			
K.1	Båre, egnet til K.5 (se note)	1 stk.	0
K.1	Stretcher, suitable for K.5 (see note)	1 pc.	0
K.2	Oppustelig frakturskinne, senior, hel arm	1 stk.	0
K.2	Inflatable fracture splint, senior, full arm	1 pc.	0
K.3	Oppustelig frakturskinne, senior, hel ben	1 stk.	0
K.3	Inflatable fracture splint, senior, full leg	1 pc.	0
K.4	Stiv halskrave, indstillelig til forskellige positioner	2 stk.	0
K.4	Stiff neck collar, adjustable	2 pcs.	0
K.5	Langt rygbræt, inkl. seler (se note)	1 stk.	0
K.5	Spineboard, incl straps see note)	1 pc.	0
K.2 - K.4 pakkes og opbevares sammen. / K.2-K.4 packed and stored together			
K.1 + K.5 Meddelelser A kapitel IX B, bilag 3 / Notice A Chapter IX B, annex 3			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A1
L. Tandlægeudstyr / Equipment for dental examination and treatment (Se note / see note)			
L.1	Coltosol®, færdigblandet tandcement i tuber à 8g (se note)	5 stk.	0
L.1	Coltosol®, dental filling material ready for use, tubes of 8 g (see note)	5 pcs.	0
L.2	Skyllekanyle	1 stk.	0
L.2	Dental rinse needle	1 pc.	0
L.3	Specialinstrument til tandcement	1 stk.	0
L.3	Special instrument for dental filling material	1 pc.	0
L.4	Mundspejl	1 stk.	0
L.4	Mouth mirror	1 pc.	0
L.5	Ekskavator	1 stk.	0
L.5	Excavator	1 pc.	0
L.6	Fyldningsfjerner	1 stk.	0
L.6	Filling remover	1 pc.	0
L.7	Rustfri stålæske med låg til tandudlægeudstyr	1 stk.	0
L.7	Instrument box, stainless steel with lid for dental equipment	1 pc.	0
L: Tandlægeudstyr skal ikke forefindes på skibe med fart indenfor 175 sømil fra EU/EØS landes kyster. L: Dental equipment shall not be on ships in voyage within 175 nautical miles from EU/EEA countries' coast L.1: Tandcement skal være inklusive instrument til uddrivning af cement fra tuben. L.1: Filling material shall include instrument for expulsion of the material from the tube.			
M. Udstyr til sygepleje / Requisites for nursing			
M.1	Medicinnålebægre, ca. 25 ml, éngangs	20 stk.	0
M.1	Graduated measuring cup, approx. 25 ml, disposable	20 pcs.	0
M.2	Varmedunk	1 stk.	0
M.2	Hot-water bottle	1 pc.	0
M.3	Bækken	1 stk.	0
M.3	Bedpan	1 pc.	0
N. Insektbekæmpelsesmidler og kloreringsmidler til rensning af drikkevand/ Insecticides and chlorinating agents for purifying drinking water (se note / see note)			
N.1	Insektbekæmpelsesmiddel - pulver form	1 dåse	0
N.1	Insecticide, not specified - powder form	1 tin	0
N.2	Kloreringsmiddel til drikkevand, 100 g	2 stk.	0
N.2	Chlorinating agent for drinking water, 100 g	2 pcs.	0
N.3	Insektbekæmpelsesmiddel- væskeform	1 dåse	0
N.3	Insecticide - liquid form	1 Tin	0
N.1 + N.2 + N.3 Klor og insektbekæmpelsesmidler må ikke opbevares sammen med lægemidler og medicinsk udstyr. N.1 + N.2 + N.3 Chlor and insecticides must not be stored within the ship medicine chest			
O. Fortegnelser, lægebog m.v. / Inventories, Medical Guide, etc.			
O.2	Lægebog for Søfarende, udgivet af Seahealth (se note)	1 stk.	0
O.2	Medical guide for seafarers issued by Seahealth (see note)	1 pc.	0
O.3	Fortegnelse, kontroldokument og brugervejledning (se note)	1 stk kat. A MFAG A1	0
O.3	Inventory, Control Document and User Instruction (see note)	1 pc. Cat. A MFAG A1	0
O.5	Radio Medical Optegnelser	1 stk.	0
O.5	Radio Medical Records	1 pc.	0
O.6	MFAG (Medical First Aid Guide)	0	1 stk.
O.6	MFAG (Medical First Aid Guide)	0	1 pc.
O.2 Skal også forefindes på engelsk, såfremt der om bord er påmønstret ikke dansktalende besætningsmedlemmer. / O.2 Must also be available in English, if crewmembers who don't speak Danish have signed on. O.3 skal være tilgængelig i digital eller printet udgave. O.3 must be available in digital or printed version.			

Labels

1.1 Gas / gas Medicinsk ilt / Medical Oxygen Indikation : Bevidstløse uden vejtrækning, Bevidstløse med vejtrækning, Medtaget almen tilstand. Slag mod kraniet, Større kvæstelser, Vejtrækningsbesvær Indication :Unconscious, Weak general condition, Blows to the skull. Severe injuries, Difficulty breathing. Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring: Opbevares adskilt fra brandbare gasser og stoffer. Hold flasketemperaturen under 50°C / Storage: Keep away from flammable gases and substances. Keep bottle under 50°C
2.1 Inj. væske / inj. liquid Lidocain 10 mg/ml Indikation / Indication: Lokalbedøvelse / Local anaesthetic Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
2.2 Gel / gel Lidocain 2% Indikation/Indication: Lokalbedøvelse / Local anaesthetic Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
3.1 Tablet / tablet Paracetamol 500 mg Indikation / Indication: Svage smerter / Weak pain; MFAG tables 7, 8,13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
3.2 Tablet / tablet Ibuprofen 400 mg Indikation / Indication: Moderate smerter / Moderate pain Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
3.4 Inj.væske / inj.liquid ▲ Morfin 10 mg/ ml Indikation / Indication: Stærke smerter / Severe pain; MFAG tables 7,8, 10,13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
3.7 Gel / ointment Diclofenac 11.6 mg/g Indikation / Indication: Smerte / pain , inflammation I huden / inflammation of the skin. Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 30 C
3.8 Supp. / supp. Diclofenac 50 mg Indikation / Indication: Galdesten, nyresten, ledsmerter / Gall stone, kidney stone, pain in joint Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
3.9 Tablet/tablets Morphine hydroclorid 10 mg Indikation / Indication: Stærke smerter / Severe pain; Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
4.1 Inj.væske / inj.liquid Adrenalin 1 mg/ml Indikation / Indication: Allergisk Shock / Allergic shock Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring: Læs indlgssedel/ Storage: Read packet insert

4.2 Kapsel / capsule Acrivastin 8 mg Indikation / Indication: Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
4.3 Inj.væske / inj.liquid Hydrocortisonsuccinat 100 mg Indikation / Indication: Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
4.6 Tablet / Tablets Prednisolon 25 mg Indikation / Indication Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
4.7 Tablet / tablet ▲ Cyclizine 50 mg Indikation / Indication: Søsyge / Seasickness Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
5.1 Inj.væske / inj.liquid Naloxone 0,4 mg/ml Indikation/Indication: Morfin antidot / Morphine antidote MFAG tables 4, 13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.3 Inj.væske / inj.liquid Atropin 1 mg/ml Indikation / Indication: Antidot / Antidote / MFAG table 17 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.4 Gel / gel Calcium gluconate 2 % Indikation / Indication: Antidot / Antidote / MFAG table 8, 16 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.5 Tablet / tablet Calciumlactogluconate 500 mg Indikation / Indication: Antidot / Antidote / MFAG tables 16 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.6 Afdelt pulver / powder Aktivt kul / Activated charcoal Indikation / Indication: Antidot / Antidote/ MFAG table 10 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.7 Væske / liquid Ethyl alcohol 99,5% Indikation/Indication: Antidot / Antidote/ MFAG table 19 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
5.8 Inj.væske / inj.liquid Phytomenadion 10 mg/ml Indikation/Indication: Antidot / Antidote / MFAG table 14 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

6.2 Tablet / tablet Δ Diazepam 5 mg Indikation / Indication: Angst / Anxiety Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
6.3 Klysma / enema Δ Diazepam 4 mg/ml Indikation / Indication: Kramper / Convulsions; MFAG tables 4, 5, 6 Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
6.4 Inj.væske / inj liquid Δ Haloperidol 5 mg/ml Indikation / Indication: Sindssyge / Mental disease Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
6.5 Tablet / tablet Δ Zolpidem 10 mg Indikation / Indication: Søvnløshed / insomnia Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.1 Tablet / tablet Phenoxymethylpenicillin 660 - 800 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.2 Tablet / tablet Amoxicillin 500 mg Indikation / Indication: Infektioner / Infections / MFAG table 9 Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.6 Tablet / tablet Metronidazol 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.7 Suppositorie / suppository Metronidazol 500 mg Indikation / Indication: Infektioner / Infections / MFAG table 10 Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.8 Tablet / tablet Ciprofloxacin 250 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.9 Inj.substans. / inj.substance Ceftriaxon 1 g Indikation / Indication: Infektioner / Infections / MFAG table 10 Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.10 Vaccine / vaccine Indeholdende / containing Tetanus Indikation / Indication: Forebyggelse mod stivkrampe / Tetanus prevention Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2- 8°C

7.11 Immunglobulin / immunglobuline Tetanus 250 I.E./ml / 250 I.U/ml Indikation / Indication: Forebyggelse mod stivkrampe / Tetanus prevention Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2- 8°C
7.12 Tablet / tablet Flucloxacillin 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.15 Tablet / Tablets Azithromycin 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.16 Tablet / Tablets Pivmecillinam 400 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
7.17 Tablet / Tablets Aciclovir 400 mg Indikation / Indication: virus-Infektioner / virus-Infections Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
8.1 Tablet / tablet Atovaquon/Proguanil 250/100 mg Indikation / Indication: Malaria forebyggelse / Malaria Prevention. Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
8.2 Tablet / tablet Doxycyklin 100 mg Indikation / Indication: Malaria forebyggelse / Malaria prevention Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
8.4 tablet / tablet Artemether and lumefantrine 20 mg / 120 mg Indikation / Indication: Malaria behandling / Malaria treatment Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: ikke over 30° C / Not above 30° C
9.1 Spray / aerosol Glycerilnitrat / Glyceril trinitrate 0,4 mg/doses Indikation / Indication: Hjertekrampe / Angina pectoris Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2 - 25°C
9.3 Tablet / tablet Acetylsalicylsyre / acetylsalicylic acid 150 mg Indikation / Indication: Blodfortyndende / Blood thinning Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
9.4 Tablet / tablet Furosemid 40 mg Indikation / Indication: Vanddrivende / Diuretic Ref: Se Fortegnelseens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

9.5 Inj.væske / inj.liquid Furosemid 10 mg/ ml Indikation / Indication: Vanddrivende / Diuretic MFAG tables 2, 9 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory. Opbevaring / storage: normal temperatur / normal temperature
9.6 Tablet / tablet Atenolol 50 mg Indikation / Indication: Blodtryksforhøjelse / High blood pressure Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
9.8 tablet / Tablets Apixaban 5 mg Indikation / Indication:Blodfortyndende/ Blood diluting Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
9.9 inj. / inj. Tranexamsyre / Tranexamacid 100 mg/ml Indikation / Indication:Stopper blødninger/ Stops Bleedings Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
10.2 Creme / lotion Miconazol 2% Indikation / Indication: Hudsvamp / Skinfungus Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 25°C
10.5 Creme / lotion Permethrin 5% Indikation / Indication: Fladlus og fnat / Crab lice and scabies Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
10.6 Shampoo / shampoo Permethrin 1% Indikation / Indication: Lus/ Lice Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
10.7 Creme / lotion Glukokortikoid gruppe II /glucocorticoid group II Indikation / Indication: Eksem / Eczema Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
10.8 Salve / oinment Fusidin 2 % Indikation / Indication:Hud infektion / Skin infection Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
11.1 Væske / liquid Chlorhexidin / Ethanol 0,5 - 5%/70 - 90% Indikation / Indication: Desinfektion af instrumenter / Disinfection of instruments Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
11.2 Væske / liquid Chlorhexidin 0,2% Indikation / Indication: Behandling af urene sår, mundskyl / Treatment of dirty wounds, mouth rinse. Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

12.6 Tablet / tablet Loperamid 2 mg Indikation / Indication: Diarré / Diarrhea Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
12.8 Kapsler / capsule Omeprazol 20 mg Indikation / Indication: Mavesår / Stomach ulcer Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
12.7 Pulver / powder Glucose / kalium / magnesium / natrium citrat/natrium/ Glucose/potassium/magnesium/sodium citrate/sodium 30/1,1/0,11/2,1 g Indikation / Indication: Væske- og salttab / Rehydration MFAG table 8, Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature
12.9 Tablet eller oral opløsning / tablet or oral solution Antacida Indikation / Indication: Sure opstød og halsbrand / acid indigestion and heartburn Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
12.10 Rektalsalve / rectal ointment Glukokortikoid Indikation / Indication: Analkløe / Anal pruitisRef: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 25 C
12.13 Tablet / Tablets Ondasetron 4 mg Indikation / Indication: Kvalme og opkastning / Nausea and vomiting MFAG table 7, 8, 10, 13, 15, 20 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / Storage normal temperatur.
12.14 Pulver / Power Movicol® Indikation / Indication: Forstoppelse / Constipation Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
13.3 Øjendråber / eyedrops Dexamethason 1 mg/ml Indikation / Indication: Allergi / Regnbuehindebetændelse / Allergic / Inflammatory eye diseases Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
13.4 Øjendråber / eyedrops Pilocarpin 20 mg/ml Indikation / Indication: Grøn stær / Glaucoma Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
13.6 Øjensalve / eyeointment Aciclovir 30 mg/g Indikation / Indication: Virusinfektion / Viral infection Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

13.9 Væske / liquid Medium til at blødgøre ørevoks/ Medium to soften earwax Indikation / Indication: Blødgøring og løsning af ørevoks / Softening and dissolving of earwax Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
13.11 Øjendråber / eyedrops Tropicamid 1% Indikation / Indication: Øjeninflammation / Inflammation of eye Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
13.12 Øredråber / eardrops Ciprofloxacin / Hydrocortison 10+2 mg Indikation: Øregangs eksem / Ear canal eczema Ref: Se Fortegnensens brugervejledning / Read user instruction in the inventory. Søg samråd med Radio Medical / Consult Radio Medical
13.13 Øjensalve / eye ointment Chloramphenicol 1% Indikation: øjen infektion /Eye infection Ref: Se Fortegnensens brugervejledning / Read user instruction in the inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: Se indlægsseddel / See leaflet
13.14 Øjendråber / Eye drops Diclofenac 1 mg/ml Indikation: Lokalbedøvelse / Local anaesthetic. Ref: Se Fortegnensens brugervejledning / Read user instruction in the inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: Ikke over 25°C / Not above 25°C
14.1 Kapsel / capsule Fluconazol 150 mg Indikation / Indication: Skedesvamp / Vaginal fungus Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
14.2 Tablet / tablet Methylergometrin 0,125 mg Indikation / Indication: Blødning efter fødsel / Bleeding after giving birth Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
15.1 Tablet / tablet Dextromethorphan 30 mg Indikation / Indication: Hoste / Cough Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

15.5 Næsespray / dråber / nose spray / drops Xylometazolin 1 mg/ml Indikation / Indication: Forkølelse / Coon cold Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
15.7 Inhalations Spray / Inhalation Spray Salbutamol 0,1 mg Indikation / Indication: Astma / Astma – MFAG table 9 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
15.8 Inhalations spray / Inhalation spray Glucocorticoid 100-200 µg Indikation / Indication: Astma / Astma Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
16.1 Inf.væske / inf.liquid Natriumklorid / sodium chloride 9 mg/ml Indikation / Indication: Blodtab / Loss of blood; MFAG table 7 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
16.2 Inf.væske / inf.liquid Glucose og natrium / glucose and sodium 25/4,5 mg/ml Indikation / Indication: Væsketab / Rehydration Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
16.5 Inj. væske / inj.liquid Natriumklorid / sodium chloride 9 mg/ml Indikation / Indication: Væske til gennemskylning af intravenøs kanyle, sårbehandling, øjenbehandling / Fluid for flushing intravenous needle, treatment of wound, treatment of eyes Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature
16.7 Væske / Liquid Sterilt vand / Sterile water Indikation / Indication: Væske til blanding af 7.9 / Fluid to mixing 7.9 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature

INDEX

Abdominal infections	31; 32	Blodprop i hjertet	14; 39; 40
Abstinenskrampes	27	Blodprop i hjertet, forebyggelse efter	42
Acetylsalicylsyre	40	Blodtab	61
Aciclovir	36; 53; 67; 70; 84; 85; 95; 97; 109	Blood diluting	43
Acid indigestion	50	Blood poisoning	33
Acrivastin	19	Blood-thinning	40
Acrivastine	19	Blurred vision	53
Active carbon	25	Blødning, abort	57
Acute diarrhoea	32	Blødning, fødsel	57
Acute mental disease	27	Boil in the throat	31; 32
Acute mental illness	28	Bone infection	31
Acyclovir	53	Bone-ache	14
Adrenalin	18	Bones, infections	35
Aktivt kul	25	Breathing difficulties	11
Akut diarré	32	Brud	27
Akut sindssyge	27; 28	Brugervejledning Lægemiddelstoffer	10
Alcoholic withdrawal symptoms	27	Burning sensation, skin	45
Alkoholabstinenser	27	Burns	15
Allantoin	54	Calciumgluconat	23
Allergi	19	Calciumgluconate	24
Allergic asthma	19; 20	Calciumlactogluconat, calciumcarbonat.	24
Allergic face oedema	19; 20	Calciumlactogluconate, calciumcarbonate.	24
Allergic reactions, swellings etc. Oral treatment preferred.	20	Cardiospasm	40
Allergic shock	18	Caustic burns, fluorine	24
Allergisk ansigtsødem	19	Caustic burns, stomach, intestines	25
Allergisk astma	19	Ceftriaxone	33
Allergisk shock	18	Cerumen	54
Allergiske reaktioner, hævelser osv.Oral behandling	20	Chlamydia infections	35
Allergy	19; 20	Chlamydia infektioner	35
Amoxicillin.	30	Chloramphenicol	55
Amoxicilline	30	Chlorhexidin.	47
Angina pectoris	39; 42	Cholera	32
Angst	27; 28	Ciprofloxacin	32
Antacid	50	Ciprofloxacin	32
Antacida	50	Cold	59
Antidote, morphine	21	Constipation	51
Antihæmorrhoid creme	50	Convulsions	27
Anxiety	27; 28	Convulsions , universal	28
Apixaban	43; 109	Convulsions, poisoning	28
Appendicitis	31; 32; 33	Convulsions, Withdrawal	28
Artemether and lumefantrine	38	Coronary thrombosis	15; 39; 40
Artemether og lumefantrine	38	Coronary thrombosis, Preventiv treatment after	42
Arthritis	15	Cough	58
asthma	59	Crab lice	44; 45
Asthma	60	Cyclizine	21
asthma-like breathing	60	Delusions	28
astma	59	Dexamethason.	52
Astma	42; 60	Dexamethasone.	52
astmalignende vejrtrækning	60	Dextromethorphan.	58
Atenolol	42	Diarré	48
Atovaquon/Proguanil	37	Diarrhoea	48; 49
Atovaquone/Proguanil	37	Diazepam	27; 28
Atropin	23	Diclofenac	15; 16; 56
Azithromycin	35; 109	Difficulty falling asleep	29
Back pains	13	Dirty wounds, Disinfection	47
Bevidstløs med vejrtrækning	11	Dislocations	27
Bevidstløs uden vejrtrækning	11	Doxycycline.	38
Bihulebetændelse	30; 59	Dysenteri	32
Bihulebetændelse	35	Dysentery	32
Bleeding, after birth	58	Ear eczema	52
Bleeding, miscarriage	58	Eksem	45
Blindtarmsbetændelse	31; 32; 33	Epilepsi	27
Blodforgiftning	33	Epilepsy	28
Blodfortyndende	43	Epinephrine	18
Blodfortynding	40	Ernæringsrestatning	61

Ethyl alcohol	26	Heart pain	39; 40; 42
Ethyl alkohol	26	Heartburn	50
Eye diseases, Allergic	52	helvedesild	36
Eye diseases, Inflammatory	52	Helvedesild	53
Eye infection	55	Herpes	53
Eyes, Treatment	62	Herpes simplex	36
Feber	13	Herpes zoster	36
Febertilstande	48	High blood pressure	41; 42
Female reproductions organs, infection	32	Hjertekrampe	39; 40
Fever	13	Hostestillende	58
Feverish conditions	49	Hovedpine	13
Fladlus	44; 45	Hudinfektioner	46
Flucloxacillin	35	Hudkløe	45
Fluconazol	57	Hudsvamp	44; 57
Fluconazole.	57	Hudsvie	45
Flushing, intravenous needle	62	Human tetanus immunoglobulin	34
Fnat	44; 45	Humant tetanus immunoglobulin	34
Food substitute	62	Hydrocortisonsuccinat.	19
Forbrændinger	14	Hydrocortisonsuccinate	20
Forgiftning, blodfortyndende pesticider	26	Hypoglycaemic coma	22
Forgiftning, insektgift	23	Hæmorider	50
Forgiftning, medikamenter	26	Hænder, desinficering	47
Forgiftninger med methanol, ethylene glycol	26	Høfeber	19
Forgiftninger, mave, tarm	25	Ibuprofen.	13; 14
Forhøjet blodtryk	40; 41; 42	Indsovningsbesvær	29
Forkølelse	59	Infection of female reproductions organs	31
Forstoppelse	51	Infections in connection with ulcer	35
Fortegnelse og kontroldokument over lægemiddelstoffer	64	Infections of the oral cavity	35
		Infections, oral cavity	30
Fortegnelse og Kontroldokument over medicinsk udstyr	73	Infektioner, stafylokokker	35
		Inflammation i huden	15
Fractures	14; 15; 27	Inflammation of the middle ear	35
Frontal sinusitis	59	Inflammation of the skin	15
Fungus	44	Inflammation of the throat	35
Furosemid	40; 41	Inflammation, middle ear	30
Furosemide	41	Inflammation, throat	30
Fusidin	46; 109	Inflammatoriske øjensygdomme	54
Galdesten	14	Inflammatory eye diseases	54
Gall stone	15	Injuries	15
Gennemskylning, intravenøskanyle	62	Insect bites	45
Gigt	15	Insecticides	23
Glaucoma	53	Insektbid	45
Glucagon	22	Insektgifte	23
Glucocorticoid til inhalation	60	Insomnia	29
Glucocorticoid group II	45	Instrumenter, desinficering	47
Glucose and sodium chloride.	62	Instruments, Disinfection	47
Glucose og natriumklorid.	61	Insulin shock	22
Glukokortikoid gruppe II	45	Insulinshock	22
Glyceryl trinitrate	39	Insulintilfælde	22
Glycerylnitrat	39	Intact skin, Disinfection	47
Glycocorticoid for inhalation	60	Intakt hud, desinficering	47
Gonorré	32	Intestinal infections	31; 32
Gonorrhoea	32	Inventory and Control Document of Active Pharmaceutical	
Grøn stær:	52	Ingredients	64
Haemorrhoids	50	Inventory and Control Document of Medical Equipment	73
Hallucinationer.	28	Itching, rectum	50
Hallucinations.	28	Kidney stone	15
Haloperidol	28	Kloramfenikol	55
Haloperidole.	28	Klorhexidin.	47
Halsbetændelse	30; 35	Klorhexidin/ethanol.	47
Halsbrand	50	Kløe, endetarmen	50
Halsbyld	31; 32	Kløestillende	12
Hands, Disinfection	47	Knoglebrud	13
Hay fever	19	Knogleinfektion	31; 35
Headache	13	Knoglesmerter	13

Kolera	32	MFAG table 8, 10, 11:	49
Kramper	27	MFAG table 10	25; 32; 33
Kramper, forgiftninger	27	MFAG table 14	26
Kramper, universelle	27	MFAG table 16	24
Kvalme	51	MFAG table 17	23
Kvalme og opkastninger	28	MFAG table 19	26
K-vitamin	26	MFAG table 2, 3, 5, 9, 11	11
Kvæstelser	14	MFAG table 2, 9	41
Kæbehulebetændelse	59	MFAG table 4, 13	21
Lavt blodsukker	22	MFAG table 4, 5, 6	27
Ledskred	27	MFAG table 8, 16	24
Ledsmerter	13	MFAG table 9	30; 35
Legionella	35	Miconazol	44
Lice	44; 45	Miconazole	44
Lidocain	11; 12		105; 106
Lidocaine	12	Moderate pains:	14
List when handing over expired medication to a pharmacy	107	Moderate smerter:	13
Liste ved aflevering af udløbet medicin til apotek	107	Modgift, morfin	21
Local anaesthetics	12	Morfin	14
Lokalbedøvelse	11; 12	Morphine	15
Loperamid	48	Morphine hydroclorid Δ	17
Loperamide.	48	Morphinhydroclorid Δ	17
Loss of blood	61	Motoric restlessness,	28
Loss of fluid	49	Motorisk uro	28
Loss of liquid	61	Mouth rinse	47
Loss of liquids	62	Movicol®	51; 85; 96
Low blood sugar	22	Mundhuleinfektioner	30; 35
Lung edema	41	Mundskylning	47
Lungebetændelse	30; 33; 35	Muscle pains	14
Lungebetændelse, giftige kemiske stoffer	30	Muskelsmerter	13
Lungepest	32	Nail fungus	57
Lus	44; 45	Naloxon	21
Major wounds	15; 33	Naloxone	21
Malaria, forebyggelse	37	Natriumklorid	62
Malaria, prevention	37; 38	Natriumklorid.	61
Malarone®	37	Nausea	51
Mani, vildskab	28	Nausea and vomiting	28
Mania, wildness	28	Neglesvamp	57
Mavesår	49	Nyrebækkenbetændelse	32; 33
Mavesårsinfektioner	35	Nyresten	14
Maxillary sinusitis	59	Nældefeber	19
Medical oxygen	11	Omeprazol	49
Medicinsk ilt.	11	Ondansetron	51; 96
Medtaget almen tilstand	11	Open fractures	33
Mellemørebetændelse	30; 35; 59	Opkastninge	51
Meningitis	33	Opkastninger	48
Menstruationssmerter	13	Otitis media	59
Mental restlessness	28	Oxygen	11
Methylergometrin.	57	P. Falciparum malaria	38
Methylergometrine.	58	P. Falciperum malaria	38
Metronidazol.	31; 32	Pain	15
Metronidazole	31	Pain during defecation	50
MFAG tabel 4, 5, 6	28	Pain in joints	14
MFAG tabel 8, 10, 11	48	Pains, eye	53
MFAG tabel 10	25; 31; 33	Pandehulebetændelse	59
MFAG tabel 14	26	Paracetamol	13
MFAG tabel 16	24	Paracetamol.	13
MFAG tabel 17	23	Period pains	14
MFAG tabel 19	26	Permethrin.	44; 45
MFAG tabel 4, 13	21	Phenoxyethylpencillin	30
MFAG tabel 7	61	Phenoxyethylpenicillin	30
MFAG tabel 7, 8, 10, 13	14	Phytomenadion	26
MFAG tabel 7, 8, 10, 13	15	Phytomenadione	26
MFAG tabel 8, 16	23	Pilocarpin	52
MFAG tabel 9	30; 35	Pilocarpine	53

Pivmecillinam	36; 109	Syning af sår	11
Pneumonia	30; 33; 35	Søsyge	21
Pneumonia, poisonous chemical substances	30	Søvnløshed	29
Pneumonic plague	32	Sårbehandling.	62
Poisoning, blood- thinning pesticides,	26	Tandbyld	31; 32
Poisoning, insecticides	23	Tarminfektioner	31
Poisoning, medicaments	26	Tetanus antidote	34
Poisoning, methanole, ethylene glycole	26	Tetanus vaccine	33; 34
Poisoning, stomach, intestines	25	tetanustoxoid	34
Prednisolon	20	Tooth abscess	31; 32
Prenisolon	20	Tranexamacid	43
Psoriasis	45	Tranexamsyre	43
Psykisk uro	28	Tropicamid	54
Pyelitis	32	Tågesyn	52
Rainbow vision	53	Udsættende vejtrækning	21
Regnbuesyn	52	Unconscious with breathing	11
Relieves itching	12	Unconscious without breathing	11
Restlessness	27	Underlivsinfektioner	31
Rygsmarter	13	Urene sår, desinficering	47
Salbutamol	59	Urethritis	32
Salbutamol.	59	Urinary tract infection	36
Salttab	48	Urinrørsbetændelse	32
Scabies	44; 45	Urinvejsinfektioner	36
Seasickness	21	uroilstande	27
Sepsis	33	Urticaria	19
Severe bronchitis	32	User instruction Active Pharmaceutical Ingredients	10
Severe injuries	11	Vaginal fungus	57
Severe pain	15; 17	Vand i lungerne	40; 41
Shingles	53	Vand til opløsning af 7.9 Ceftriaxon pulver	63
Shock, forebyggelse	61	Variceller	36
Shock, prevention	61	Vejtrækningsbesvær	11
Sinusitis	30; 35; 59	vitamin K	26
Skin fungus	44; 57	Vomiting	49; 51
Skininfections	46	Vrangforestillinger	28
Skoldkopper	36	Væsketab	48; 61
Smerter	15	Water for dissolution of 7.9 Ceftriaxon powder	63
Smerter i hjertet	39; 40	Weak general condition	11
Smerter, afføring	50	Weak pains:	13
Smerter, øje	52	Wounds, Treatment	62
Sodium chloride	61; 62	Xylometazolin.	59
Solskoldning	45	Xylometazoline.	59
Staphylococci	35	Zolpidem	29
Sterile water	63	Ændringer /Changes	111
Sterilt vand	63	Ændringer/ Changes	110
Stivkrampemodgift	34	Ætsninger, fluorholdige	23; 24
Stivkrampevaccine	33	Ætsninger, mave, tarm	25
Stomach ulcer	49	Øjenbehandling	62
Stopper blødninger	43	Øjenbetændelse	55
Stops Bleedings	43	Øjensygdom, Allergiske	52
Stops breathing	21	Øjensygdom, Inflammatoriske	52
Store sårskader	14; 33	Øregangseksem	52
Stærke smerter	17	Ørevoks.	54
Stærke smerter:	14	Åbne knoglebrud	33
Større kvæstelser	11		
Sunburns	45		
Sure opstød	50		
Suture of wounds	12		
Svage smerter	13		
Svamp	44		
Svamp i skeden	57		
Svær bronkitis	32		

**Rekvisition lægemiddelstoffer og medicinsk
udstyr**

**Requisition Active Pharmaceutical Ingredient
and medical equipment**

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
1.1	V03AN01	Medicinsk ilt	Gas	
1.1	V03AN01	Oxygen, Medical	Gas	
1.3	-	Ekstra iltflaske	Gas	
1.3	-	Spare oxygen cylinder	Gas	
2.1	N01BB02	Lidocain	Inj. Væske	
2.1	N01BB02	Lidocaine	Inj. Liquid	
2.2	D04AB01	Lidocain	Gel	
2.2	D04AB01	Lidocaine	Gel	
3.1	N02BE01	Paracetamol	Tablet	
3.1	N02BE01	Paracetamol	Tablets	
3.2	M01AE01	Ibuprofen	Tablet	
3.2	M01AE01	Ibuprofen	Tablets	
3.4	N02AA01	Morfin ^Δ	Inj. Væske	
3.4	N02AA01	Morphine ^Δ	Inj. Liquid	
3.7	M02AA15	Diclofenac	Gel	
3.7	M02AA15	Diclofenac	Gel	
3.8	M01AB05	Diclofenac	Supp.	
3.8	M01AB05	Diclofenac	Supp	
3.9	N02AA01	Morphinhydroclorid ^Δ	Tablet	
3.9	N02AA01	Morphine hydroclorid ^Δ	Tablets	
4.1	C01CA24	Adrenalin "K"	inj. Væske	
4.1	C01CA24	Epinephrine "R"	inj. Liquid	
4.2	R06AX18	Acrivastine	Kapsel	
4.2	R06AX18	Acrivastine	Capsula	
4.3	H02AB09	Hydrocortison	inj. tørstof væske	
4.3	H02AB09	Hydrocortisonsuccinate	inj. Subst + solvens	
4.6	H02AB06	Prednisolon	Tablet	
4.6	H02AB06	Prednisolon	Tablets	
4.7	R06AE03	Cyclizin ^Δ	Tablet	
4.7	R06AE03	Cyclizine ^Δ	Tablets	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
5.1	V03AB15	Naloxon	Inj. Væske	
5.1	V03AB15	Naloxon	Inj. Liquid	
5.2	H04AA01	Glucagon "K"	inj. tørstof + væske	
5.2	H04AA01	Glucagon "R"	inj. Subst + solvens	
5.3	A03BA01	Atropine	Inj. Væske	
5.3	A03BA01	Atropine	Inj. Liquid	
5.4	D11AX03	Calcium gluconate	gel	
5.4	D11AX03	Calcium gluconate	gel	
5.5	A12AA02 / A12AA06	Calcium lactogluconate	Brusetablet	
5.5	A12AA02 / A12AA06	Calcium lactogluconate	Effervescent tablet	
5.6	A07BA01	Aktivt kul	pulver	
5.6	A07BA01	Charcoal, activated	powder	
5.7	-	Ethyl alcohol	Væske	
5.7	-	Ethyl alcohol	Liquid	
5.8	B02BA01	Phytomenadione	Inj. Væske	
5.8	B02BA01	Phytomenadione	Inj. Liquid	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
6.2	N05BA01	Diazepam Δ	Tablet	
6.2	N05BA01	Diazepam Δ	Tablets	
6.3	N05BA01	Diazepam Δ	Klysma	
6.3	N05BA01	Diazepam Δ	Enema	
6.4	N05AD01	Haloperidol Δ	Inj. Væske	
6.4	N05AD01	Haloperidol Δ	Inj. Liquid	
6.5	N05CF02	Zolpidem Δ	Tablet	
6.5	N05CF02	Zolpidem Δ	Tablets	
7.1	J01CE02	Phenoxymethylpenicillin	Tablet	
7.1	J01CE02	Phenoxymethylpenicillin	Tablets	
7.2	J01CA04	Amoxicillin	Tablet	
7.2	J01CA04	Amoxicillin	Tablets	
7.6	P01AB01	Metronidazol	Tablet	
7.6	P01AB01	Metronidazol	Tablets	
7.7	P01AB01	Metronidazol	Supp.	
7.7	P01AB01	Metronidazol	Supp.	
7.8	J01MA02	Ciprofloxacin	Tablet	
7.8	J01MA02	Ciprofloxacin	Tablets	
7.9	J01DD04	Ceftriaxon	inj. tørstof	
7.9	J01DD04	Ceftriaxon	inj. Subst	
7.10	J07AM01	Tetanus holdig vaccine "K"	Inj. Væske	
7.10	J07AM01	Vaccine containing Tetanus "R"	Inj. Liquid	
7.11	J06BB02	Tetanus immunglobulin "K"	Inj. Væske	
7.11	J06BB02	Tetanus immunglobulin "R"	Inj. Liquid	
7.12	J01CF05	Flucloxacillin	Tablet	
7.12	J01CF05	Flucloxacillin	Tablets	
7.15	J01FA10	Azithromycin	Tablet	
7.15	J01FA10	Azithromycin	Tablets	
7.16	J01CA08	Pivmecillinam	Tablet	
7.16	J01CA08	Pivmecillinam	Tablets	
7.17	J05AB01	Aciclovir	Tablet	
7.17	J05AB01	Aciclovir	Tablets	
8.1	P01BB51	Atovaquon/Proguanil	Tablet	
8.1	P01BB51	Atovaquon/Proguanil	Tablets	
8.2	J01AA02	Doxycycline	Tablet	
8.2	J01AA02	Doxycycline	Tablets	
8.4	P01BF01	Artemether and lumefantrine	Tablets	
8.4	P01BF01	Artemether and lumefantrine	Tablets	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
9.1	C01DA02	Glyceryl nitrat	Mund pray	
9.1	C01DA02	Glyceryl trinitrate	Oral spray	
9.3	B01AC06	Acetylsalisylsyre	Tablet	
9.3	B01AC06	Acetylsalicylic Acid	Tablets	
9.4	C03CA01	Furosemid	Tablet	
9.4	C03CA01	Furosemide	Tablets	
9.5	C03CA01	Furosemide	Inj. væske	
9.5	C03CA01	Furosemide	Inj. Liquid	
9.6	C07AB03	Atenolol	Tablet	
9.6	C07AB03	Atenolol	Tablets	
9.8	B01AF02	Apixaban	Tablet	
9.8	B01AF02	Apixaban	Tablets	
9.9(π)	B02AA02	Tranexamsyre	Inj. væske	
9.9(π)	B02AA02	Tranexamsyre	Inj. liquid	
10.2	D01AC02	Miconazol	Crema	
10.2	D01AC02	Miconazol	Lotion / Cream	
10.5	P03AC04	Permethrin	Crema	
10.5	P03AC04	Permethrin	Lotion / Cream	
10.6	P03AC04	Permethrin	Shampoo	
10.6	P03AC04	Permethrin	Shampoo	
10.7	D07AB02	Glukocorticoid group II	Crema	
10.7	D07AB02	Glukocorticoid group II	Lotion / Cream	
10.8	D06AX01	Fusidin	Cream / Salve	
10.8	D06AX01	Fusidine	Cream / Lotion	
11.1	-	Chlorhexidine/ethanol	Væske	
11.1	-	Chlorhexidine/ethanol	Liquid	
11.2	-	Chlorhexidine	Væske	
11.2	-	Chlorhexidine	Liquid	
12.6	A07DA03	Loperamid	Tablet	
12.6	A07DA03	Loperamid	Tablet	
12.7	-	Glucose/potassium/magnesium /sodium	Powder	
12.7	-	Glucose/potassium/magnesium /sodium	Powder	
12.8	A02BC01	Omeprazol	Tablet / Kapsel	
12.8	A02BC01	Omeprazol	Tablet / Capsula	
12.9	A02A	Antacida	Tablet eller oral opløsning	
12.9	A02A	Antacid	Tablet or oral solution	
12.10	C05A	Hæmoride creme(skal indeholde glukokortikoid and smertestillende)	Crema/Salve	
12.10	C05A	Anti haemorrhoid agent (Must contain glucocorticoid and analgesi)	Lotion/ointment	
12.13	A04AA01	Ondansetron	Tablet	
12.13	A04AA01	Ondansetron	Tablets	
12.14	A06AD65	Movicol®	Pulver	
12.14	A06AD65	Movicol®	Powder	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
13.3	S01BA01	Dexamethason	Dråber	
13.3	S01BA01	Dexamethasone	Drops	
13.4	S01EB51	Pilocarpin	Dråber	
13.4	S01EB51	Pilocarpine	Drops	
13.6	S01AD03	Aciclovir	Salve	
13.6	S01AD03	Aciclovir	Ointment	
13.9	-	Medium til at blødgøre ørevoks, inkl. Øresprøjte	-	
13.9	-	Medium to soften earwax, incl. Earsyringe	-	
13.11	S01FA06	Tropicamid	Dråber	
13.11	S01FA06	Tropicamide	Drops	
13.12	S02CA03	Ciprofloxacin / Hydrocortison	øretråber	
13.12	S02CA03	Ciprofloxacin / Hydrocortisone	eardrops	
13.13	S01AA01	Kloramfenikol	øjensalve	
13.13	S01AA01	Chloramphenicol	eye ointment	
13.14	S01BC03	Diclofenac	Øjendråber	
13.14	S01BC03	Diclofenac	Eye drops	
14..1	J02AC01	Fluconazol	Kapsel	
14.1	J02AC01	Fluconazole	Capsula	
14.2	G02AB01	Methylergometrin	Tablet	
14.2	G02AB01	Methylergometrine	Tablets	
15.1	R05DA09	Dextromethorphan	Tablet	
15.1	R05DA09	Dextromethorphone	Tablets	
15.5	R01AA07	Xylometazolin	Dråber / spray	
15.5	R01AA07	Xylometazoline	Drops / spray	
15.7	R03AC02	Salbutamol	Spray	
15.7	R03AC02	Salbutamol	Spray	
15.8	R03BA01	Glucocorticoid (inhalation)	Spray	
15.8	R03BA01	Glucocorticoid (inhalation)	Spray	
16.1	B05BB01	Natriumklorid	Inf. Væske	
16.1	B05BB01	Sodium chloride	Inf. Liquid	
16.2	B05BB02	Glucose + Natriumklorid	Inf. Væske	
16.2	B05BB02	Glucose + sodiumchlorid	Inf. Liquid	
16.5	V07AB	Natriumklorid 20 ml	Væske	
16.5	V07AB	Sodium chloride 20 ml	Liquid	
16.7	V07AB	Sterilt vand til blanding af 7.9	Væske	
16.7	V07AB	Sterile water to mix 7.9	Kiquid	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A
A.1	Mobilt genoplivningsudstyr, klargjort	
A.1	Mobile resuscitation equipment, ready for use	
A.4	Tungeholdere, 80 mm	
A.4	Guidel airway, 80 mm	
A.5	Tungeholdere, 90 mm	
A.5	Guidel airway, 90 mm	
A.6	Tungeholdere, 100 mm	
A.6	Guidel airway, 100 mm	
A.7	Ekstra ventilationspose med ventil og voksenmaske	
A.7	Extra ventilation bag with valve and mask	
A.8	Kraftig klædesaks	
A.8	Strong scissors for clothes	
A.9	Sugekatetre, lige, CH 12	
A.9	Suction catheters, straight, CH 12	
A.10	Tæppe til behandling af hypotermi	
A.10	Blanket for treatment of hypothermia	
A.11	Almindelige tæpper	
A.11	Ordinary blankets	
A.12	Saturationsmåler (fingermodel)	
A.12	Puls oximeter (finger)	
A.13	Hudson maske med reservoirpose med ca. 1,8 m slange	
A.13	Hudson mask with reservoir bag with approx. 1.8 m tube	
A.18	Hjertestarter (AED)	
A.18	Automatic extern difibrillator (AED)	
A.19	Iltbrille	
A.19	Nasal Cannula	
B.1	Sterile gazeekompresser, ca.5x5 cm, ca.8 lag	
B.1	Sterile gauze compress, approx.5x5 cm, approx. 8 layers	
B.2	Sterile gazeekompresser, ca. 10 x 10 cm, ca. 8 lag	
B.2	Sterile gauze compress, approx.10x10 cm, approx. 8 layers	
B.3	Steril absorberende ikke-klæbende forbinding, ca. 20 x 30 cm	
B.3	Sterile absorbent non-adhesive surgical dressingbandage, approx. 20 x 30 cm	
B.4	Sterile kompresforbindinger, ca. 8 cm	
B.4	Sterile compress bandages, approx. 8 cm	
B.5	Imprægnerede net, ca. 10 x 10 cm, Jelonet	
B.5	Impregnated nets, approx. 10 x 10 cm, Jelonet	
B.6	Sterile absorberende forbindinger, ikke klæbende, ca. 10 x 10 cm	
B.6	Sterile absorbent bandages, non-adhesive surgical dressingbandages, approx. 10 x 10 cm	
B.7	Gazeviskers, usterile	
B.7	Gauze swaps, non-sterile	
B.8	Tubegaze til fingerforbindinger	
B.8	Tube shaped gauze for finger bandaging	
B.9	Elastisk gazebind, ca. 4 cmx4m	
B.9	Elastic gauze bandage, approx. 4 cmx4m	
B.10	Hæfteplaster med sårpude, forskellige størrelser, lommepakning	
B.10	Adhesive tape with compress, various sizes, pocket pack	
B.11	Hæfteplaster med sårpude, ca. 6 cmx1m	
B.11	Adhesive tape with compress, approx. 6 cmx1m	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quentety
B.12	Fikseringsplaster Fixomull, ca. 10 cm x 10 m	
B.12	Adhesive tape Fixomull, approx. 10 cm x 10 m	
B.13	Fikseringsplaster, "Englehud", ca. 2 cm	
B.13	Adhesive tape, "peau d'ange", approx. 2 cm	
B.18	Årepresse til massiv blødning	
B.18	Tourniquet for massive bleeding	
C.1	Engangshandsker af latex, usterile, X-large	
C.1	Disposable gloves, latex, non-sterile, X-large	
C.2	Engangshandsker af latex, usterile, large	
C.2	Disposable gloves, latex, non-sterile, large	
C.5	Vaskesvampe, éngangs	
C.5	Bath sponge, disposable	
C.6	Beskyttelsesbriller	
C.6	Protective goggles	
C.7	Steril afdækning, ca. 50 x 50 cm	
C.7	Sterile drape, approx. 50 x 50 cm	
C.8	Flydende medicinsk sæbe pH 4,5	
C.8	Liquid medical soap pH 4,5	
C.9	Mundbind 3 lags, bredde 18 cm	
C.9	Mask, 3 layer, width 18 cm	
C.10	Ligpose voksen	
C.10	Body bag adult	
C.11	Engangshandsker af latex, usterile, medium	
C.11	Disposable gloves, latex, non-sterile, medium	
C.12	Engangshandsker af latex, usterile, small	
C.12	Disposable gloves, latex, non-sterile, small	
C.13	Engangsforklæde	
C.13	Disposable apron	
C.14	Lagen, sterilt	
C.14	Bed sheet, sterile	
C.15	Absorberende underlag, ca.6 lag, 60x40 cm, usteril, engangs	
C.15	Absorbent underlay, approx.6 layers, 60x40 cm, non-sterile, disposable	
C.16	Opkastpose	
C.16	Vormit bag	
D.1	Suturtape, Steri-Strip™	
D.1	Suturtape, Steri-Strip™	
D.2	Klæbesutur, Dermabond	
D.2	Adhesive suture, Dermabond	
D.4	Ikke-absorberbar sutur, 3.0; 45 cm. FS eller DS 25	
D.4	Non-absorbable suture, 3.0; 45 cm. FS or DS25	
D.5	Rustfri stålæske med låg, ca. 18x9x3 cm	
D.5	Instrument box, stainless steel with lid, approx. 18x9x3 cm	
D.6	Engangsskalpel, steril, nr. 23	
D.6	Disposable scalpel, sterile, no. 23	
D.7	Kirurgisk saks med lige spids	
D.7	Surgical scissors w/straight points	
D.8	Kirurgisk saks med krum spids	
D.8	Surgical scissors w/curved points	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quentety
D.9	Kirurgisk pincet, ca. 14 - 16 cm	
D.9	Surgical forceps (tweezers), approx. 14 -16 cm	
D.10	Anatomisk pincet	
D.10	Anatomical forceps (tweezers)	
D.11	Splintpincet	
D.11	Forceps, splinter (tweezers)	
D.12	Pean, krum	
D.12	Pean, curved	
D.13	Nåleholder	
D.13	Suture - holder	
D.14	Barbermaskine (skraber), engangs	
D.14	Safety razor, disposable single use	
D.15	Absorberbar hæmostatisk gelatinesvamp, ca.1x1x1 cm	
D.15	Haemostatic absorbable gelatine sponge, approx.1x1x1 cm	
D.17	ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25	
D.17	Non-absorbable suture, 5.0; 45 cm. FS eller DS25	
E.1	Kanylebøtte	
E.1	Box for contaminated needles	
E.2	Afspritningssvaps m. isopropylalkohol 70 %	
E.2	Alcohol swaps, isopropyl alcohol 70 %	
E.3	Staserem	
E.3	Tourniquet for i.v.	
E.4	Intravenøs kanyle, 1,1 x 32, 20 g	
E.4	Intravenous needle, 1.1 x 32, 20 g	
E.5	Fikseringsplaster til intravenøs kanyle	
E.5	Fixing plaster for intraveneos needle	
E.6	Kanyle, steril til engangsbrug, 0,6 x 30 (~ 23G x 1 1/4)	
E.6	Needle, sterile, disposablesingle use, 0.6 x 30 (~ 23G x 1 1/4)	
E.7	Sprøjter, sterile til engangsbrug, Luer, 2 ml	
E.7	Syringes, sterile, single use, Luer, 2 ml	
E.8	Sprøjter, sterile til engangsbrug, Luer, 5 ml	
E.8	Syringes, sterile, single use, Luer, 5 ml	
E.9	Sprøjter, sterile til engangsbrug, Luer, 10 ml	
E.9	Syringes, sterile, single use, Luer, 10 ml	
E.10	Infusionssæt med gevindtilslutning til E.4, sterile til engangsbrug	
E.10	Infusion set with threaded connection for E.4, sterile, single use	
F.1	Blodtryksmåler	
F.1	Instrument for measuring blood pressure	
F.2	Stetoskop	
F.2	Stethoscope	
F.3	Digitalt lægetermometer	
F.3	Digital clinical thermometer	
F.4	Pencillygte m. batterier	
F.4	Pencil lamp w/batteries	
F.5	Tungespatel, usterile, engangs	
F.5	Tongue depressor, non-sterile, single use disposable	
F.7	Hypotermi Termometer	
F.7	Hypotermi thermometer	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quentety
G.1	Multistix® 5, urintest for blod/glukose/protein/nitrit/leukocytter	
G.1	Multistix® 5, Urinetest for blood/glucose/protein/nitrite/leukocytes	
G.2	Graviditetstest	
G.2	Pregnancy test kit	
G.5	Malaria test, minimum for Plasmodium Falciparum, inkl. reagens og brugervejledning.	
G.5	Malaria test, minimum for Plasmodium Falciparum, incl. reagens and userinstruction.	
G.6	Lancetter (fingerprikker)	
G.6	Lancets (finger pricker)	
G.7	Udstyr til måling af blodsukker	
G.7	Equipment for measuering bloodsugar	
G.8	Hurtig test C-reaktiv protein (CRP)	
G.8	Rapid test for C- reaktive proteine (CRP)	
H.1	Pencillygte med UV lys, inkl. batterier	
H.1	Lamp with UV light w/batteries	
H.2	Fluoretsstrips til farvning af øjet	
H.2	Fluorescent colour for the eye	
H.3	Vatpinde, sterile	
H.3	Cotton sticks, sterile	
H.5	Øjenklap	
H.5	Eye patch	
H.6	Øjenskylllevæske, Natriumchlorid 0,9%, 250 ml	
H.6	Eye irrigating liquid, Sodium chlorid e 0.9%, 250 ml	
I.1	Engangskatetre m. krum spids, CH 14, til mænd	
I.1	Catheters for single use w/ curved tip, CH 14 for men	
I.2	Urinkelbe, plastic	
I.2	Bed bottle, plastic	
I.3	Engangskatetre CH 14 til kvinder	
I.3	Catheters for single use, CH 14 for women	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quentety
J.1	ColdHot™ kulde/varme omslag, ca.11 x 27 cm	
J.1	ColdHot™ pack approx. 11 x 27 cm	
J.2	Elastikbind, ca. 10 cm x 5m	
J.2	Elastic bandage, approx. 10 cm x 5m	
J.3	Elastikbind, ca. 6 cm x 5m	
J.3	Elastic bandage, approx. 6 cm x 5m	
J.4	Trekantet tørklæde (Mitella)	
J.4	Triangular scarf (Mitella)	
J.5	Sikkerhedsnåle	
J.5	Safety pins	
J.6	Fingerskinne til immobilisering af ét led	
J.6	Finger splint for immobilisation of one joint	
J.7	Fingerskinne til immobilisering af to led	
J.7	Finger splint for immobilisation of two joints	
J.8	Polsterbind, ca. 10 cm x 2,7 m	
J.8	Padded bandages, approx. 10 cm x 2.7 m	
J.9	Gipsbind, ca. 12 cmx2m	
J.9	Plaster of paris bandage, approx. 12 cmx2m	
J.10	Fikseringsplaster, Tensoplast, ca. 2 cmx2m	
J.10	Adhesive tape, Tensoplast, approx. 2 cmx2m	
K.1	Båre, egnet til K.5	
K.1	Stretcher, suitable for K.5	
K.2	Oppustelig frakturskinne, senior, hel arm	
K.2	Inflatable fracture splint, senior, full arm	
K.3	Oppustelig frakturskinne, senior, hel ben	
K.3	Inflatable fracture splint, senior, full leg	
K.4	Stiv halskrave, indstillelig til forskellige positioner	
K.4	Stiff neck collar, adjustable	
K.5	Langt rygbræt, inkl. seler	
K.5	Spineboard, incl straps	
L.1	Coltosol®, færdigblandet tandcement i tuber à 8g	
L.1	Coltosol®, dental filling material ready for use, tubes of 8 g	
L.2	Skyllekanyle	
L.2	Dental rinse needle	
L.3	Specialinstrument til tandcement	
L.3	Special instrument for dental filling material	
L.4	Mundspejl	
L.4	Mouth mirror	
L.5	Ekskavator	
L.5	Excavator	
L.6	Fyldningsfjerner	
L.6	Filling remover	
L.7	Rustfri stålæske med låg til tandudlægeudstyr	
L.7	Instrument box, stainless steel with lid for dental equipment	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quentety
M.1	Medicinmålebægre, ca. 25 ml, éngangs	
M.1	Graduated measuring cup, approx. 25 ml, disposable	
M.2	Varmedunk	
M.2	Hot-water bottle	
M.3	Bækken	
M.3	Bedpan	
N.1	Insektbekæmpelsesmiddel - pulver form	
N.1	Insecticide, not specified - powder form	
N.2	Kloreringsmiddel til drikkevand, 100 g	
N.2	Chlorinating agent for drinking water, 100 g	
N.3	Insektbekæmpelsesmiddel- væskeform	
N.3	Insecticide - liquid form	
O.2	Lægebog for Søfarende, udgivet af Seahealth (se note)	
O.2	Medical guide for Seafarers issued by Seahealth	
O.3	Fortegnelse, kontroldokument og brugervejledning kat A MFAG A1(Download, www.soefartsstyrelsen)	
O.3	Inventory, Control document and User instruction Cat A MFAG A1 (Download, www.dma.dk)	
O.5	Radio Medical Optegnelser	
O.5	Radio Medical Records	
O.6	MFAG (Medical First Aid Guide)	
O.6	MFAG (Medical First Aid Guide)	
O.2 Skal også forefindes på engelsk, såfremt der om bord er påmønstret ikke dansktalende besætningsmedlemmer. O.2 Must also be available in English, if crewmembers who don't speak Danish have signed on. O.3 skal være tilgængelig i digital eller printet udgave. www.soefartsstyrelsen.dk O.3 must be available in digital or printed version. www.dma.dk		

Mobilt genoplivnings udstyr / Akut taske
Mobile resuscitation kit / Emergency bag

Mobilt genoplivnings udstyr		Version 1-2024
Lovpligtige udstyr.		
Grp. No.	Medicin - Medicinsk udstyr	Mængde
1.3	O ₂ flaske, 2-4 l med 200 Bar,	1 stk.
A.5	Tungeholder, 90 mm	1 stk.
A.7	Ventilationspose med voksen maske og tilhørende slange	1 stk.
A.20	Mekanisk sug med kateter	1 stk.
A.21	Reduktionsventil/Flowregulator	1 stk.
O.8	HLR ombord DK/UK(blå A4)	1 stk.
O.9	Instruktion Reduktionsventil/Flowregulator, DK/UK	1 stk.
Forslag til medicin og udstyr (ikke lovpligtigt)		
A.8	Klædesaks	1 stk.
A.10	Tæppe til behandling af Hypotermi	1 stk.
A.13	O ₂ Hudson maske med reservoirpose	1 stk.
A.19	Iltbrille	1 stk.
A.22	Gaffa tape	1 rulle
B.2	Sterilt gaze kompres, ca.10 x 10 cm	2 stk.
B.3	Absorberende forbinding, ca. 20 x 30 cm	2 stk.
B.4	Steril gaze forbinding, med kompresfelt ca. 8 cm	2 stk.
B.6	Absorberende forbinding, ca.10 x 10 cm	2 stk.
B.9	Elastisk gazebind (ikke sterilt) ca. 4 cm x 4 m	2 stk.
B.13	Rulle tape, hudvenlig, ca. 2 cm bred	1 stk.
B.18	Årepresse / Tourniquet til massiv blødning	1 stk.
C.2	Handsker L, ikke-sterile	6 par
E.1	Kanylebøtte	1 stk.
E.2	Afspritningsswabs med isopropyl 70 % alkohol	10 stk.
E.3	Staserem (tourniquet til iv)	1 stk.
E.4	Venekanyle	4 stk.
E.5	Fikserings plaster til iv kanyler	4 stk.
E.6	Kanyle til injektion	4 stk.
E.7	2 ml sprøjte	4 stk.
E.10	Infusion sæt (drop-slange)	1 stk.
F.1	Blodtryksapparat	1 stk.
F.2	Stetoskop	1 stk.
F.4	Pencillygte	1 stk.
K.2	Oppustelig skinne (Urias), senior, hel arm	1 stk.
K.3	Oppustelig skinne (Urias), senior, helt ben	1 stk.
K.4	Halskrave, justérbar (ANLÆGGES KUN PÅ ORDINATION FRA RADIO MEDICAL)	1 stk.
K.6	Mekanisk pumpe til K.2 og K3 (ikke obligatorisk om bord)	1 stk.
O.5	Radio Medical rapport DK eller UK	1 stk.
O.10	Indholdsfortegnelse (har du i hånden)	1 stk.
O.11	Kuglepen	1 stk.
O.12	Halskrave Flowchart – ny instruks	1 stk.
O.13	Halskrave Actioncard– ny instruks	1 stk.
O.14	Clipboard	1 stk.
O.15	Aktionskort, ABCDE, DK eller UK	1 stk.
9.1	Nitroglycerin, spray m 200 doser	1 stk.
11.2	Flaske Klorhexidin 0,2 %	1 stk.
16.1	Saltvand (NaCl) infusionsvæske	1 stk.
16.5	Saltvand (NaCl) 20 ml plastik flaske	2 stk.

Mobile resuscitation equipment		Version 1-2024
Equipment according to Legislation.		
Grp. No.	Medicine - Medical equipment	Quantity
1.3	O ₂ bottle, 2-4 l with 200 bar	1 pc
A.5	Guedel Airway, 90 mm	1 pc.
A.7	Ventilation bag with adult mask and associate tubing	1 pc
A.20	Mechanical suction with catheter	1 pc
A.21	Reduction valve/Flowregulator	1 pc
O.8	CPR onboard (Blue A4) DK/UK	1 pc
O.9	Instruction Reduction valve/Flowregulator DK/UK	1 pc
Suggestions for medicine and equipment (not legally required)		
A.8	Pair of scissors for clothes	1 pc
A.10	Blanket for treatment of hypothermia	1 pc
A.13	O ₂ Hudson mask with reservoir bag	1 pc
A.19	Nasal Cannula	1 pc
A.22	Duch tape	1 role
B.2	Sterile gauze compress, approx. 10 x 10 cm	2 pcs.
B.3	Absorbing bandage, approx. 20 x 30 cm	2 pcs.
B.4	Sterile, elastic gauze, with compress approx. 8 cm	2 pcs.
B.6	Absorbing bandage, approx. 10 x 10 cm	2 pcs.
B.9	Elastic gauze, (non-sterile) approx. 4 cm x 4 m	2 pcs.
B.13	Roll of adhesive tape for skin, approx. 2 cm wide	1 pc
B.18	Tourniquet for massive bleeding	1 pc
C.2	Gloves L, non-sterile	6 pair
E.1	Needle bin	1 pc
E.2	Alcohol swap with isopropyl 70 % alcohol	10 pcs.
E.3	Tourniquet for IV	1 pc
E.4	Intravenous Cannula	4 pcs.
E.5	Fixating cover, for IV needles	4 pcs.
E.6	Needle for injection	4 pcs.
E.7	2 ml syringe	4 pcs.
E.10	Infusion set (hose)	1 pc
F.1	Instrument for measuring blood pressure	1 pc
F.2	Stethoscope	1 pc
F.4	Pen light	1 pc
K.2	Inflatable splint (Urias), whole arm	1 pc
K.3	Inflatable splint (Urias), senior whole leg	1 pc
K.4	Stiff neck collar, adjustable (ONLY APPLIED IF PRESCRIBED BY RADIO MEDICAL)	1 pc
K.6	Mechanical pump for K.2 and K.3 <i>/not mandatory onboard</i>	1 pc
O.5	Radio Medical Record	1 pc
O.10	List of Contents (in your hand)	1 pc
O.11	Pen	1 pc.
O.12	Neck collar Flowchart - new guide	1 pc.
O.13	Neck collar Actioncard - new guide	1 pc.
O.14	Clipboard	1 pc.
O.15	Action card ABCDE DK/UK	1 pc
9.1	Glyceryl trinitrate , spray 200 dosis	1 pc.
11.2	Bottle Chlorhexidine 0.2	1 pc.
16.1	Sodium chloride (NaCl) Inf. Liquid 1 liter	1 pc.
16.5	Sodium chloride (NaCl) 20 ml	2 pcs.

Liste ved aflevering af udløbet medicin til apotek
List when handing expired medication to a pharmacy

Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
Morfin ^Δ	Inj. Væske	
Morphine ^Δ	Inj. Liquid	
Morphine hydroclorid ^Δ	Tablet / Kapsel	
Morphine hydrochloride ^Δ	Tablet / Capsula	
Cyclizin ^Δ	Tablet	
Cyclizine ^Δ	Tablets	
Diazepam ^Δ	Tablet	
Diazepam ^Δ	Tablets	
Diazepam ^Δ	Klysma	
Diazepam ^Δ	Enema	
Haloperidol ^Δ	Inj. Væske	
Haloperidole ^Δ	Inj. Liquid	
Zolpidem ^Δ	Tablet	
Zolpidem ^Δ	Tablets	
Skib - kontrol nr. Ship - control no.	Skibsførerens Navn og Underskrift Masters Name and signature	
Dato / Date	Apotek / Pharmacy, eller/or Agent	
Navn / Name	Underskrift / Signature	

Ændringer / Changes

Version 3.5 July 2025

A.4 – 6 tungeholder farve kode og størrelse fjernet – **mm** angivelsen er mest sikker.

12.14 Korrekte lægemiddelstof erstatter "Movicol® "

15.7 Inhalation Salbutamol ATC kode rettet i liste over medicin.

15.7 Ligeledes er holdbarheden rette i brugervejledningen

Label 4.6 tilføjet " *Søg samråd med Radio Medical / Consult Radio Medical*"

A.4 – A.6 Guidal airway: Color coding and size removed – the millimeter specification is the most reliable.

12.14 The correct active substance replaces "Movicol®".

15.7 Inhalation Salbutamol: ATC code corrected in the list of medicines.

15.7 The validity has also been corrected in the user manual.

Label 4.6 added: "*Consult Radio Medical / Søg samråd med Radio Medical.*"

October version 3.4 2024

Tilføjet side med liste over lovpligtig udstyr i mobilt genoplivningsudstyr og forslag til ekstra indhold

3.9 Morphin er præciseret til Morphinhydrochlorid

13.12 **Ciprofloxacin / Hydrocortison** tilføjet i July 2024 opdateringen - manglede i Ændringer/Changes.

Liste til aflevering af udløbet medicin til apotek er rette.

A.4 – A.6 Tungeholder er nu defineret i størrelse med farve og mm

Added page with list of mandatory equipment in mobile resuscitation equipment and suggestions for additional content

3.9 Morphine is specified as Morphine hydrochloride

13.12 **Ciprofloxacin / Hydrocortisone** added in the July 2024 update - was missing from the Ændringer/Changes.

-List to deliver expired medicine to pharmacy is corrected.

A.4 – A.6 Guidel airway is now defined in size with color and mm.

August 2024

4.5 Cinnarizin er udgået af markedet. Erstatte af 4.7 Cyclizin **Δ**

4.5 Cinnarizine is withdraw from market. It is replaced with 4.7 Cyclizine. **Δ**

July-2024

13.2 udgået af markedet

13.14 Diclofenac erstatter 13.2.

13.13 Kloramphenicol Opbevaring ændret til:: Se indlægsseddel

13.2 Withdrawn from market.

13.14 Diclofenac replaces 13.2

13.13 Chloramphenicol Storage changed to: See leaflet

July-2024

13.8 Fusidin øjensalve er nu fjernet fra medicinliste

13.8 Fusidin eye ointment is now removed from list of medicine.

Medicin som er fjernet / Medicine removed

3.5 Tbl. Tramadol 50 mg
7.4 Tbl. Erythromycin 500 mg
7.13 Tbl. Sulfamethoxazol+trimitoprim
8.3 Tbl. Quinin 100 mg
10.4 Cream Sølvulfadiazin
12.5 Mixtur Laktulose 667 mg/ml
12.12 Tbl. Dompemridon 10 mg
13.8 Dråber/gel Fusidin
15.2 Turbohaler Terbutalin
15.4 Turbohaler Budesonide
15.6 Inj. Salbutamol
16.6 Gelatine, Natriumchlorid

Medicin som er tilføjet / Medicine added

3.9 Tbl. Morfin 10 mg
4.6 Tbl. Prednisolon 25 mg
7.2 Tbl. Amoxicillin 500 mg
7.15 Tbl. Azithromycin 500 mg
7.16 Tbl. Pivmecillinam 400 mg
7.17 Tbl. Aciclovir 400 mg
8.4 Tbl. Artemether and
9.8 Tbl. Apixaban 5 mg
9.9 Inj. Tranexamsyre 100 mg/ml
10.8 Salve / Lotion Fusidin 2%
12.13 Tbl. Ondansetron 4 mg
12.14 Pulver / Powder Movicol®
13.12 Ciprofloxacin / Hydrocortisone
13.13 Øjensalve / ointment Chloramphenicol
15.7 Inhalation spray Sabumetol
15.8 Inhalation spray glyocortocoid
16.7 Steril vand / Sterile water

Medicinsk udstyr som er fjernet / Medical equipment that is removed

E.4 Optiva*2 Intravenøs kanyle, 1,1 x 32, 20 g / Optiva*2 intravenous cannula, 1.1 x 32, 20 g
E.5 Vecafix® Fikseringsplaster til intravenøs kanyle / Vecafix® Fixing plaster for intravenous cannula

Medicinsk udstyr som er tilføjet / Medical equipment that is added

A.18 Hjertestarter (AED) 7 / Automatic external defibrillator (AED)
A.19 Iltbrille / Nasal Cannula
C.15 Absorberende underlag ca. 6 lag 60x40 cm, usteril, engangs / Absorbent underlay, approx. 6 layers, 60x40 cm, non-sterile, disposable
C.16 Opkastpose / Vomitbag
D. 17 Ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25 / Non-absorbable suture, 5.0; 45 cm. FS eller DS25
E.4 Intravenøs kanyle, 1,1 x 32, 20 g (produkt navn fjernes) / intravenous cannula, 1.1 x 32, 20 g (production name removed)

E.5 Fikseringsplaster til intravenøs kanyale (produkt navn fjernes) / Fixing plaster for intraveneos cannula (product name removed)
G.8 Hurtig test C-reaktiv protein (CRP) / Rapid test for C- reaktive proteine (CRP)
I.1 Engangskatetre m. krum spids, CH 14, til mænd (antal øges til 10 stk.) / Catheters for single use w/ curved tip, CH 14 for men (the amount increases to 10 pcs.)

October 2023

4.1 Adrenalin opbevaring ændret - Læs indlægssedlen for opbevaring
16.3 Hydroxyethylstivelse er fjernet fra medicin oversigt.
4.5 Cinnarizin π fjernet da det er håndkøb
E.4 og E.5 Engelsk tekst rettet fra "needle" til cannula.
Tilføjet liste aflevering af udløbet medicin på apotek
Generelle rettelser af stavfejl og layout

4.1 Epinephrine storage changed - read packet insert for storage
4.5 Cinnarizin π removed - is an over the counter purchase
16.3 Hydroxyethyl amoid remove from list of medicine.
E.4 og E.5 Text corrected from "needle" to cannula.
Added list of delivery of expired medicine at the pharmacy
General spelling corrections and layout

April 2022

B.14 rettet til B.18 Årepresse massiv blødninger
Generelle rettelser af stavfejl og layout.

B.14 corrected to B.18 Tourniquet for massive bleeding
E.3 correted to Tourniquet **for i.v.**
General spelling corrections and layout.

October 2021

Tilføjet Identifikation af skib (Side 1)
Tilføjet 13.12 Ciprofloxacin / Hydrocortison- øredråber
16.3 Hydroxyethylstivelse udgår
Tilføjet 16.6 Gelatine, Natrium chlorid erstatter 16.3
Tilføjet B.14 Årepresse til pulsåreblødning
Tilføjet C.13 Engangsforklæde
Tilføjet C.14 Sterilt lagen
Tilføjet F.7 Hypotermi termometer
Tilføjet I.3 Engangskatetre CH 14 til kvinder
Ændret N.1 Insektbekæmpelsesmiddel, ikke specificeret- **pulver form**
Tilføjet N.3 Insektbekæmpelsesmiddel, ikke specificeret- **væskeform**

Added Identification of ship (page 1)
Added 13.12 Ciprofloxacin / Hydrocortisone - eardrops
16.3 Hydroxyethyl amoid is discontinued product
Added 16.6 Gelatine sodium Chlorid replace 16.3
Added B.14 Tourniquet to arterial bleeding
Added C.13 Disposable apron
Added C.14 Bed sheet, sterile
Added F.7 Hypotermi thermometer
Added I.3 Catheters for single use, CH 14 for women
Changed N.1 Insecticide, not specified - **powder form**

July 2021

6.3 Diazepam klysma styrke rettet tilbage til 4mg/ml - beklager ulejligheden

7.10 Vaccine Tetanus- label tilrettet

COVID-19 tillæg - overskrifter rettet. Tilføjet at hurtigtest skal være CE mærket

6.3 Diazepam enema strength corrected back to 4mg/ml - Sorry for the inconvenience

7.10 Vaccine Tetanus - label corrected

COVID-19 add on - the headlines corrected. Added that the quick test have to be CE marked.

February 2021

Tilføjet 8.4 Artemether og lumefantrine i "add medicine covid-19". Til behandling af Malaria.

Add. 8.4 Artemether and lumefantrine in "add medicine covid-19". For treatment of Malaria.

January 2021

8.2 Doxycyclin styrke ændret til 100 mg

8.2 Doxycyclin strength changed to 100 mg

Tilføjet side med ekstramedicin og medicinsk udstyr i forbindelse med Covid-19. 102-> Det er frivilligt om skibet vil medbringe medicinen og udstyret eller dele deraf.

Added page with additional medicine and medical equipment in connection with Covid-19. 102-> It is optional if the ship will bring the medicine and equipment or part of it.

October 2020

6.3 Klysma Diazepam styrke ændret til 5 mg/ml

6.3 Enema Diazepam strength changed to 5 mg/ml

7.10 Tetanus holdig vaccine - styrke fjernet, da det kommer an på hvad der udleveres - følg råd fra Radio Medical

7.10 Tetanus-containing vaccine - strength removed as it depends on what is delivered - follow advice from Radio Medical

Rettet i 8.1 Brugervejledning for behandlingen af Malaria

Corrected 8.1 User instruction for treatment of Malaria

Marts 2020

Ændringer er nu på sidste side

Changes is now at the last page

15.3 Inj. Terbutalin 0,5 mg/ml er erstattet med 15.6 Inj. Salbutamol 0,5 mg/ml

15.3 Inj. Terbutalin 0,5 mg/ml is changed to 15.6 Inj. Salbutamol 0,5 mg/ml

7.1 Phenoxymethylpenicillin styrken er ændret til 660 mg - 800 mg

7.1 Phenoxymethylpenicillin strenght change to 660 mg - 800 mg

D.4 Sutur fjernet firma navn.

D.4 Sutur removed company name

August 2019

3.6 ændret til: 3.8 Diclofenac suppositorier 50 mg 50 stk.

7.10 ændret til tetanustoksid holdig vaccine 1. stk.

8.1 (Behandling) Brugervejledning rettet

O.2 Tilføjet ny lægebog for søfarende fra Seahealth

3.6 Changed to: 3.8 Diclofenac suppositorie 50 mg 50 pcs.

7.10 Changed to: Vaccine containing Tetanustoxid 1 pc.

8.1 (Treatment) User instruction corrected.

